




2




3




4



5



6



7


Pain

- ~ Pain is very important to our survival
- ~ Pain is defined as the perception of a noxious (harmful) stimulus
- ~ Pain can also occur in the absence of injury or long after an injury has healed
- ~ Pain provides humans with information about:
 - * Tissue-damaging stimuli
 - * Thus enables them to protect themselves from greater damage
- ~ Pain is protective in two ways:
 - * It removes a person from stimuli that cause tissue damage through withdrawal reflexes
 - * Learning associated with pain causes the person to avoid stimuli that previously caused pain
- ~ Pain often initiates the search for medical assistance and helps us to pinpoint the underlying cause of disease

8

Somatosensory System

- ~ Diverse sensory system composed of the receptors and processing centers to produce the sensory modalities:
 - * Touch
 - * Temperature
 - * Proprioception (body position)
 - * Nociception (pain)
- ~ The system reacts to diverse stimuli using different receptors
 - * Thermoreceptors
 - * Nociceptors
 - * Mechanoreceptors
 - * Chemoreceptors



9

Pain

- ~ Pain is an unpleasant sensory experience associated with actual or potential damage to the body, or perception of such damage. It is a subjective experience
- ~ Subjective experience
- ~ Memories of events associated with extreme pain persist for a long time
- ~ Mental state is known to have a powerful influence over pain
 - * An athlete may not notice a twisted ankle until after the competition is over.
 - * Soldiers in battle often continue to fight even after sustaining serious injury, and they may report afterwards that they experienced no pain until after battle
- ~ The scientific explanation for this phenomenon is that the brain not only receives pain messages, but also has a descending system of neurons that suppresses pain messages

10


Pharmacology of Pain Management

- ~ **Peripheral acting agents**
 - * Prevent sensitization of receptors to substance P
 - * Example: NSAIDs, ibuprofen
- ~ **Signal inhibiting agents**
 - * Prevent pain signal from travelling to cortex
 - * Example: Anesthetics, proparacaine
- ~ **Central acting agents**
 - * Act on pain perception centers in the cortex (CNS)
 - * Example: opioids/narcotics

11


Descending Pathway

- ~ This system inhibits cells in the spinal cord that transmit pain signals
- ~ A pathway for natural pain modulation
- ~ Opioids that occur naturally such as the endorphins are important neurotransmitters in some of these descending pathways



12

Peripheral versus Central Acting



13

Four Major Types of Pain

- ☞ Nociceptive Pain
 - * Typically the result of tissue injury
- ☞ Inflammatory Pain
 - * An abnormal inflammation caused by an inappropriate response by the body's immune system
- ☞ Neuropathic Pain
 - * Pain caused by nerve irritation
- ☞ Functional Pain
 - * Pain without obvious origin but can cause pain

14

Drug Treatment Options... Neuropathic Pain

- ☞ Not the focus of today's discussion...
- ☞ Why is this relevant?
- ☞ Adjuvants – means “add on” medications
 - * Some of them have addiction potential
 - ☐ Anti-seizure medications that address nerve damage/inflammation
 - MOA: work on the GABA system – similar to benzodiazepines (ex. Xanax)
 - Gabapentin (Neurontin) – controlled substance in multiple states
 - Pregabalin (Lyrica) – controlled substance in all 50 states
 - ☐ Anti-anxiety and sleep medications
 - Zolpidem (Ambien)
 - Alprazolam (Xanax), Lorazepam (Ativan), Diazepam (Valium)

15

Neuropathic Pain Chronic Pain

- ☞ Trigeminal neuralgia
- ☞ Post-herpetic neuralgia
- ☞ Diabetic neuropathy
- ☞ Phantom limb pain following an amputation
- ☞ Multiple sclerosis
- ☞ Pain following chemotherapy
- ☞ HIV infection
- ☞ Alcoholism
- ☞ Tension headache
- ☞ Migraine
- ☞ Fibromyalgia
- ☞ Low back pain

- ☞ Tricyclic antidepressants for pain
 - * The most effective type of antidepressant used for pain
 - * Imipramine Tofranil
 - * Clomipramine Anafranil
 - * Nortriptyline Pamelor
 - * Desipramine Norpramin
- ☞ Anticonvulsants for pain
 - * Gabapentin Neurontin
 - * Topiramate Topamax
 - * Pregabalin Lyrica
 - * Carbamazepine Tegretol
 - * Oxcarbazepine Trileptal

16

Acute versus Chronic Pain

- ☞ Acute
 - * Where we are most of the time as optometrists
 - * Acetaminophen
 - * NSAIDS
 - * Opioid
- ☞ Chronic
 - * Acetaminophen
 - * NSAIDS
 - * Opioid
 - * Tricyclic antidepressants
 - * Gabapentin (Neurontin)

17

Goals of Pain DO Differ...

The goal for managing **acute pain** is to keep the patient as comfortable as possible while minimizing the **adverse drug reactions (ADRs)** from the pain meds.


The goals for managing **chronic pain** are to keep the patient as comfortable as possible (this may not mean the patient is pain free) and integrating the patient back into a “normal life” and activities of daily living, while minimizing the ADRs from the pain meds.

18

Pain Assessments and Scales

- ☞ Adds objective data to a patient's feeling of pain
 - * It is a subjective problem to assess!
 - * Remember...no patient should needlessly suffer!
- ☞ “Does the injury or wound or diagnosis fit the patient's presentation?”
 - * It is important to be able to assess the degree of pain in a patient.

19



20



21

Drug Treatment Options... Nociceptive Pain

Groups of analgesics

- * Non-opioids
 - Acetaminophen (Tylenol)
 - NSAIDs (Ibuprofen, naproxen sodium)
 - Glucocorticosteroids (methylprednisolone, prednisone)
- * Opioids
 - Codeine (Tylenol with codeine)
 - Hydrocodone (Vicodin)
 - Tramadol (Ultram)

22

Controlled Substance Schedules

Schedule I – not considered to be medically necessary, research only

- * Heroin
- * “Medical” Marijuana
 - State control of marijuana and CBD
- * LSD
- * Mushrooms
- * Ecstasy

Schedule II – **more likely to be abused (as compared to Schedule III, IV, V)**

- * Opioids, AKA “Narcotics”
 - Oxycodone (OxyContin)
 - Hydrocodone (Vicodin, Lorcet, Norco)
 - Morphine (MSContin, MSIR)
 - Hydromorphone (Dilaudid)
 - Methadone
 - Fentanyl (Duragesic)
- * ADD/ADHD meds:
 - Methylphenidate (Ritalin)
 - Mixed amphetamine salts (Adderall)

23

Controlled Substance Schedules

Schedule III – Safer, less likely to be abused (as compared to Schedule II)

- * Combination products with APAP or ASA (codeine)
- * Esketamine – nasal spray for treatment resistant depression

Schedule IV – Safer, less likely to be abused (as compared to Schedule II and III)

- * Tramadol (Ultram)
- * Benzodiazepines (lorazepam, diazepam, oxazepam)
- * Sleep agents (zolpidem, etc.)

Schedule V – safest, least likely to be abused

- * Expectorants with codeine

24

State-By-State Restriction

- ☞ Marijuana
 - * Still considered to be “C1” or “Schedule I”
 - * Federal government “ignores” it
- ☞ Hydrocodone products
 - * C3 to C2 as of 2014
 - * “hydrocodone exception”
 - NJ, etc.

25

Opioids “narcotics”

- ⌚ Mainstay of therapy for the treatment of pain
- ⌚ NO maximum daily dose limitation
- ⌚ Useful for acute and chronic pain
- ⌚ They mimic the actions of endogenous opioid compounds:
 - * Enkephalins, dynorphins, endorphins

26

Mechanisms of Action

- * Relieve pain and induce euphoria by binding to the opioid receptors (mu, kappa, delta) in the brain and spinal cord:
 - **Mu, kappa, delta** receptors in other places = ADRs
 - Mu: analgesia, **euphoria**, miosis, sedation, constipation, respiratory depression, addiction
 - Kappa: analgesia, diuresis, sedation, miosis, **dysphoria**, psychomimetic effects, respiratory depression, constipation
 - Delta: analgesia

27

Opioid Receptor Class	Effects
Mu	Analgesia, respiratory analgesia, miosis, addiction, euphoria, bradycardia, constipation
Kappa	Analgesia, diuresis, sedation, miosis, dysphoria, psychomimetic effects, respiratory depression, constipation
Delta	Analgesia, respiratory analgesia, miosis, sedation, constipation, respiratory depression, constipation

Mu, Delta, and Kappa Receptors

28

OPIOIDS IN THE BODY

29


	mu (μ, MOP or OP3)	delta (δ, DOP or OP2)	kappa (κ, KOP or OP1)
Endogenous peptides			
Beta-endorphin	+	+++	-
Leu-enkephalin	++	+++	-
Met-enkephalin	++	+	+++
Dynorphin	++		
Opioid drugs			
Pure agonists			
Morphine, codeine, oxycodone, dextropropoxyphene, methadone	+++	+	+
Pethidine, fentanyl, sufentanil	+++	-	-
Etorphine, buprenorphine	+++	+++	+++
Partial/inverse agonists			
Pentazocine, ketocyclazocine	X	+	++
Nalbuphine	X	+	(++)
Naloxone	XX	-	(++)
Buprenorphine	(+++)	-	XX
Antagonists			
Naloxone	XXX	X	XX
Naltrexone, diprenorphine	XXX	X	XXX

X: agonist activity; (+): partial agonist activity; X: antagonist activity; -: weak or no activity

30

Opioid Drugs That Bind to Opioid Receptors

- ⌚ **Opiates**
 - * Morphine, codeine
- ⌚ **Semi-synthetic**
 - * Oxycodone, hydrocodone
 - * Naloxone, Naltrexone
- ⌚ **Synthetic**
 - * Non-morphinians
 - Fentanyl
 - Methadone
 - Tramadol



31

Formulations

- ~ **Immediate release**
 - * AKA short-acting
 - * Uses: acute pain
 - Percocet, Tylenol w/ codeine, tramadol, Vicodin
- ~ **Controlled release:**
 - * AKA long-acting; sustained release; extended release
 - * Uses: basal control of chronic pain
 - * Typically NOT for acute pain nor in opioid naive patients!
 - OxyContin, MS Contin, Duragesic patch

32

Morphine Products

- ~ Standard for comparison of other agents
- ~ Used for severe pain
- ~ Multiple Brand/Trade names for long-acting morphine products, with very diverse delivery and release systems
 - * **MSIR** (IR caps) (q 3-4 hours prn)
 - * **MS Contin** (CR tabs) (q 8-12 hours)
 - * **Kadian** (CR caps) (q 12 - 24 hours)
 - * **Avinza** (CR caps) (q 24 hours)

33

Hydromorphone Products

- ~ **Hydromorphone (Dilaudid)** tablets – immediate release
 - * Take 1 – 2 tablets every 4 to 6 hours as needed for pain
- ~ **Hydromorphone ER (Exalgo)** tablets – extended release
- ~ Used for severe pain
- ~ Very potent
 - * Compare to morphine
 - 30mg PO morphine = 8mg PO hydromorphone

34

Codeine-Based

- ~ **Codeine – C3; Schedule III**
 - * Naturally occurring opioid
- ~ **Hydrocodone – C2; Schedule II**
 - * Semi-synthetic derived from codeine
 - * More potent than codeine
 - * Retains cough suppression
- ~ **Oxycodone – C2; Schedule II**
 - * Semi-synthetic derived from codeine
 - * Pain only, **no cough** suppression

35

Codeine tablets

- ~ WEAK analgesic: 30mg PO morphine = 200mg PO codeine
 - * Weakest of morphine, hydrocodone, and oxycodone
- ~ Add acetaminophen/ aspirin – Schedule III
 - * Tylenol #2 = 300 mg acetaminophen & 15 mg codeine
 - * **Tylenol #3** = 300 mg acetaminophen & 30 mg codeine
 - * Tylenol #4 = 300 mg acetaminophen & 60 mg codeine
- * 1 – 2 tablets every 4 – 6 hours as needed for pain
 - Not to exceed **3 grams** of APAP per day
- ~ Add expectorant – Schedule V

36

Oxycodone Products

- ~ **Long-Acting, Extended-Release**
 - * OxyContin
- ~ **Immediate Release; short-acting tablets**
 - * OxyIR (IR cap)
 - * Roxicodone solution
- ~ Combination with acetaminophen
 - * Percocet and Endocet (oxycodone/APAP dose)
- ~ Take 1 – 2 tablets by mouth every 4 to 6 hours as needed for pain
 - * Not to exceed 3 grams of APAP per day

37

Oxycodone Products

- ⚡ **Percodan** (oxy + asa) – no one uses this product
- ⚡ **Percocet**
 - * Oxycodone is combination with acetaminophen
 - * Various strengths
- ⚡ 30mg PO morphine = 20mg PO oxycodone

38

Hydrocodone Products

- ⚡ As of August 2014, hydrocodone products are ALL CII
 - * Moved from schedule III to schedule II
- ⚡ Immediate-Release Products
 - * **Hydrocodone 7.5 mg + IBU 200 mg**
 - **Vicoprofen**
 - * **Hydrocodone + acetaminophen:**
 - **Vicodin** = 5/300, 7.5/300, 10/300
 - **Lortab** = 2.5/300, 5/300, 7.5/300, 10/300
 - **Norco** = 5/325, 7.5/325, 10/325
- ⚡ Take 1 – 2 tabs/caps every 4 – 6 hours as needed for pain
 - * Not to exceed 3 grams of APAP per day
- ⚡ 30mg PO morphine = 20mg PO hydrocodone

39

Tramadol – another great choice

Tramadol (Ultram) tabs
Tramadol with 325 mg APAP (Ultracet), Tramadol ER tabs

- ⚡ tramadol (50 – 100 mg q 4 – 6 hours; do not exceed 400 mg/day)
- * Dual action: **mu** receptors & inhibits neuronal uptake of **serotonin** & **norepinephrine**
- * Lowers seizure threshold; increases serotonin levels
 - Watch drug interactions with other meds that ↑ serotonin
 - Selective serotonin reuptake inhibitors (SSRIs): fluoxetine/Prozac
 - Migraine meds (“triptans”): sumatriptan/Imitrex
- * **Not controlled**
 - AS OF AUGUST 2014, NOW A C4 (Schedule IV)
 - “tramies” = abuse potential; helps decrease withdrawal symptoms

40

Miscellaneous

- ⚡ **Fentanyl Patch (Duragesic)**
 - * **MOST potent opioid**
 - * **Black Box Warning** against use in acute pain and in opioid naïve patients
- ⚡ **Meperidine (Demerol)**
 - * **ACTIVE metabolites = undesirable**
- ⚡ **Methadone**
 - * Typically reserved for morphine/codeine allergic patients

41

Methadone tidbits...

- ⚡ Chronic pain or opioid abuse deterrent
- ⚡ 2-phase elimination
 - * Alpha phase = 8 hrs
 - Offers pain control
 - * Beta phase = 16+ hrs
 - Mitigates withdrawal symptoms
- ⚡ **Patient 1: On a short-acting pain med = likely being used to treat chronic pain**
 - * Twice per day dosing
- ⚡ **Patient 2: On methadone ONLY; lower doses**
 - * Once daily dosing

42


Analgesic Medications in Pregnancy

- ⚡ **Acetaminophen (Tylenol)**
 - * Analgesic of choice in pregnancy
- ⚡ **NSAIDs should generally be avoided in pregnancy**
 - * Despite Category B
 - * Miscarriage risk in first trimester
 - Ibuprofen
 - * Second trimester use is likely safe
 - Ibuprofen
 - * Third trimester avoid ALL NSAIDs
 - Premature Ductus Arteriosus closure in third trimester
- ⚡ **Opioids should be avoided in pregnancy unless there is no viable alternative**
 - * First trimester use is associated with heart defects and spina bifida

43

Opioid Allergies

- ~ If a patient states "codeine allergic" ask appropriate questions
 - * "You have indicated that you have an allergy to codeine, can you describe what happens when you take codeine?"
- ~ If a patient is truly allergic to codeine
 - * Most likely allergic to morphine, hydromorphone, oxycodone, hydrocodone, and tramadol
- ~ And...if they had an opioid IV after surgery, then their "reaction" may have been due to histamine release
 - * NOT always an allergic reaction



44

Opioid Allergies

- ~ Do you know what a patient can take if true codeine allergy?
 - ☐ Fentanyl
 - ☐ Methadone
 - ☐ Meperidine
- ~ Assessing "allergies" appropriately helps practitioner sort through Actual allergy potential and "placebo allergies"
 - ☐ Fear versus drug seeking

45




46

Specific Medications Using Numeric Pain Scale

Mild pain = 1 – 3

- ~ Acetaminophen (APAP; Tylenol)
- ~ Ibuprofen (Advil, Motrin)
- ~ Naproxen sodium (Aleve)
- ~ Tramadol (Ultram) - low dose



Moderate pain = 4 – 6


- ~ Tramadol (Ultram) – mid to high dosing
- ~ Tylenol with codeine (Tylenol #3)
- ~ Acetaminophen with oxycodone (Percocet)
- ~ Acetaminophen with hydrocodone (Vicodin) – lower dosing

47

Specific Medications Using Numeric Pain Scale

Severe pain = 7 – 10

- ~ Tylenol with hydrocodone
 - * Vicodin, etc. – higher doses
- ~ Tylenol with oxycodone
 - * Percocet, etc. – higher doses
- ~ Morphine (MSIR)
- ~ Hydromorphone (Dilaudid)
- ~ Fentanyl (Duragesic patch; Actiq lozenge on a stick)



48

"Ceiling Effect"

- ~ Commonly used when discussing *analgesics*
- ~ Phenomenon in which a drug reaches a maximum effect
 - * Increasing the drug dosage does not increase its effectiveness
- ~ Central Nervous System Agents
 - * No ceiling effect
 - * **Part of the problem**
- ~ Peripheral Nervous System Agents
 - * Has a ceiling effect

49

Tolerance

- ⚡ Escalation of dose to maintain effect
 - * Analgesia or euphoria
 - * Happens to everyone
- ⚡ Regarding euphoria = may be life threatening because respiratory depression does not show much tolerance

50


Opioid Effects/ADRs

- ⚡ CONSTIPATION-anticipate it!
 - * **All** patients should receive a stool softener + stimulant
 - * Combo: docusate + senna/Senna+S
- ⚡ Sedation
- ⚡ Euphoria – mu receptors
- ⚡ Dysphoria/Hallucinations – kappa receptors
- ⚡ Pruritis – allergy versus normal release of histamine
- ⚡ Nausea/vomiting
 - * Triggers CTZ
 - * Codeine "allergy"

51

Opioid Effects/ADRs

- ⚡ Confusion
- ⚡ Miosis
- ⚡ Respiratory depression
 - * This is what kills a patient
 - * **Mixing opioids with other CNS depressants**
 - ☐ Alcohol
 - ☐ Benzodiazepines
 - ☐ Muscle relaxers
 - ☐ Sleep agents
 - ☐ Antihistamines
 - ☐ Anti-seizure medications



52

Opioid Effects/ADRs

- ⚡ Withdrawal symptoms:
 - * Short half-life agents are more likely to cause abrupt withdrawal symptoms
 - * Sweating
 - * High sympathetic tone: increase in heart rate and blood pressure, mydriasis
 - * Agitation
 - * Irritation
 - * Irrational behavior
 - * Symptoms disappear with (immediate) use of an opioid

53


Respiratory Affects

- ⚡ Inhibition of cough reflex
- ⚡ Respiratory depression
 - * This is what kills a patient
 - * **Important to make sure that the patient doesn't**
 - ☐ Increase dose on their own
 - ☐ Add another CNS depressant with it!

54

Overdosing

- ⚡ Opioid antagonists
- ⚡ **Naloxone (Narcan) & Naltrexone (ReVia)**
 - * Used to treat opioid overdose




55


Opioid Antagonist

Naloxone (Narcan) & Naltrexone (ReVia)

- * Used to treat opioid overdose



Morphine



Naloxone

56

Mixed Opioid Agonist-Antagonist For the Treatment of Abuse/Addiction

- Exhibit partial agonist or antagonist activity at the opioid receptors
- Agonist/Antagonist combinations for the treatment of opioid abuse/addiction**
 - * **Buprenorphine (Buprenex)**
 - * Buprenorphine/Naloxone (Suboxone)
- Schedule III**
- Adverse effects
 - * Less respiratory depression & less abuse potential?
 - * Precipitate withdrawal in an opioid-dependent patient

57

Mixed Opioid Agonist-Antagonist for the Treatment of Chronic Pain

- Exhibit partial agonist or antagonist activity at the opioid receptors
- Agonist/Antagonist combinations for the treatment of chronic pain**
 - * **Not appropriate for the treatment of acute pain**
 - * Morphine/Naltrexone (Embeda)
 - * Oxycodone/Naltrexone (Troxyca ER)
- Schedule II controlled substance**

58

Substance Abuse History

- Avoid all opioids in a patient with a history of heroin use
 - * This includes tramadol
 - * May trigger dopamine reward and the drug "need"
 - * Stick with higher doses of a NSAID +/- acetaminophen
- Patients with abuse history for other substances
 - * Ex. Benzodiazepines, alcohol, amphetamines?
 - * It is a judgement call
 - * Some evidence to suggest that all addictive meds should be avoided!

59

"True Addiction" formerly "Psychological Dependence"

- Compulsive use despite harm
- Quality of life is not improved by the medication and eventually it becomes compulsive
 - * "Wanting without liking"
- Relapse is very common even after "successful" withdrawal
 - * It is a relapsing disease that is incredibly hard to treat

60

Identifying Behaviors of Abuse/Addiction

- "Fast talkers"
- Strange allergies
- Excuses for "loss" of meds
- Excuses why they need "a strong pain medication"

61

Ways to respond

- ~ Avoid getting "bullied"
- ~ Avoid acting like you are judging the patient
- ~ Use the tools that are available
 - * Call your local pharmacy/pharmacist
 - * State databases
 - o PDMP = Prescription Drug Monitoring Program
- ~ Legal/ethical issues
 - * If you didn't write it down, then it didn't happen!

62


Prescription Monitoring Program (PMP)



The Oklahoma Board of Examiners in Optometry (www.okboardopt.com) has received a Bureau of Justice Assistance grant for fund state-wide integration of the Prescription Monitoring Program with Oklahoma Health Records (OHR). Integrating the PMP with OHR provides an automated alert system to providers. The integration addresses the need for providers to have to log in separately to the PMP. Instead, the OHR automatically links a patient query and returns the patient's controlled substance prescription record directly within the provider's OHR. For more information on integration please visit <http://okboardopt.com/updates>

Sincerely,
 Russel Laveris, MD
 Executive Director
 Oklahoma Board of Examiners in Optometry

63



Dear Gregory Gaskwill,

In November 2019, **Act 112 of 2019, Opioid Treatment Agreements** went into effect. This law requires prescribers in Pennsylvania to educate their patients and establish treatment agreements with their patients before issuing the first prescription in a single course of treatment for chronic pain with a controlled substance containing an opioid.

Act 112 of 2019 requires prescribers to:

- Assess whether the patient has taken or is currently taking a controlled substance medication for treatment of a substance use disorder.
- Counsel the patient on treatment goals, expectations, benefits, and risks.
- Obtain consent from the patient for targeted urine testing.
- Include the brand name or generic name, quantity and initial dose of the controlled substance medication containing an opioid being prescribed.
- Ensure the patient understands the prescriber and patient treatment responsibilities and the prescribing policies of the clinical practice.

• Ensure the patient understands that a controlled substance medication containing an opioid has a potential for abuse, the associated risks of addiction and overdose, increased risk factors of addiction, the dangers of taking a controlled substance medication containing an opioid with benzodiazepines, alcohol or other central nervous system depressants, and other information deemed appropriate by the prescriber under 21 CFR 201.37(a)(18).

• Discuss the efficacy, risks, and benefits of other treatment options, if applicable.

To all prescribers in meeting the requirements of Act 112 of 2019, the Pennsylvania Department of Health has developed [guidance](#) for prescribers and their patients. These materials include a treatment agreement checklist, questions and answers, and patient education materials. Materials are available on the [Department of Health Website](#).

Temporary Regulations for Act 112 of 2019 were submitted for publication in the PA Bulletin on March 7, 2020, subject to change.

If you have any questions about Act 112 of 2019, please email act112@pa.gov.

Sincerely,
 Pennsylvania Department of Health

64

Ways to Combat Abuse

Drug Company Approaches

65

OxyCONTin (Controlled release tablets (q 12 hours...once in a while q 8 hours): new formulation is out to help control abuse

Manual Crushing Followed by Dissolution



Crushed New Formulation **Crushed Original Formulation**

66

Tampering for IV Abuse

- New formulation results in gelatinous material which cannot be drawn into a syringe for injection (the syringe is empty)

New formulation



Original formulation



67



68




69




70



71



72



73

Acetaminophen (Tylenol)

- o Mechanism: largely unknown
- o Mild to moderate pain
- o No anti-inflammatory potential
- o Available in 325mg, 500mg, and 650mg tablets/capsules
- o Dosing: 1,000mg every 6 to 8 hours OR 650mg every 6 hours
 - o Max daily dose: DO NOT EXCEED 3,000 to 4,000mg in 24 hours
 - o OK to use ALONG with or ALTERNATING with ibuprofen or naproxen
- o ADRs: avoid in patients who consume > 3 alcoholic beverages per day

74

NSAIDs – Ibuprofen (Advil/Motrin)

- Mechanism: prostaglandin inhibitors = decrease in inflammatory mediators
- Good for pain and inflammation
- Mild to moderate pain
- Available in 200mg (OTC) and 400mg, 600mg, and 800mg tablets (RX only)
- Dosing: 200mg to 800mg every 6 to 8 hours
 - Max daily dose: do not exceed 3,200mg in 24-hour period
 - MUST reach 1,200mg daily to achieve anti-inflammatory potential

75

NSAIDs – Naproxen Sodium (Aleve)

- o Mechanism: prostaglandin inhibitors = decrease in inflammatory mediators
- o Good for pain and inflammation
- o Mild to moderate pain
- o Available in 220mg, 275mg, 375mg, and 550mg tablets
- o Dosing: 220 to 440mg every 8 to 12 hours OR 660mg every 24 hours OR 550mg every 12 hours
 - o Acute pain: more often is BETTER
 - o Maximum daily dose is 1,000 to 1,100mg in 24 hours period
 - o OK to dose 1,375mg to 1,500mg on DAY 1 ONLY!
 - o Anti-inflammatory potential: dose at HIGHER END of range

76

NSAIDs – Adverse Effects

- o Take with food – tough on the stomach
- o May cause vasoconstriction in the kidneys
- o Inhibits platelet aggregation, so ibuprofen interacts with warfarin (Coumadin) = ↑ INR
- o May increase risk of heart attack and stroke in patients at "high risk" and with "regular use"
- o May increase blood pressure and IOP

77

SYNERGY...

It is acceptable to use an ALTERNATING dosing regimen OR an ADDITIVE dosing schedule

Good in moderate to severe pain

Acetaminophen + Ibuprofen
Ibuprofen: OTC: 200mg, Rx: 400mg, 600mg, 800mg.
Acetaminophen: OTC: 325mg, 500mg, 650mg.
Two 200mg ibuprofen every four hours while awake.*
Two 325mg acetaminophen every four hours while awake.
Maximum Daily Doses: Ibuprofen: 3,200mg. Acetaminophen: 4,000mg.
Take with food. Avoid in patients who drink three or more alcoholic beverages per day.
See previous section regarding precautions with NSAIDs. Alternate ibuprofen and acetaminophen every two hours (e.g., ibuprofen at 8am, acetaminophen at 10am, ibuprofen at 12pm, acetaminophen at 2pm, etc...)

78

Pain Reliever Help

Know your maximum daily allowances:

- ~ APAP 3000 mg (4000 mg*)
- ~ ASA 6000 mg
- ~ Ibuprofen 3200 mg
- ~ Naproxen Sodium 1650 mg (Aleve/Anaprox)
- ~ Naproxen 1500 mg (Naprosyn)

2 ibuprofen and 2 Tylenol
4 ibuprofen and 2 Tylenol

79

Alternative?

Cannabis Sativa

Hemp
Higher CBD
Traces of THC (0.2-0.3%)
Medicinal uses
Various industrial uses
Food

Marijuana
Higher THC (0-25%)
Recreational uses
Medicinal uses

non-psychoactive (no high) psychoactive (gets you high)

CBD **THC**

80

History: Prohibition of Cannabinoids

- Prohibition has only been around for 80 years
 - * Widely used for 5000-8000 years before
- Not outlawed due to lack of efficacy or safety
- Outlawed due to political and money reasons
 - * 1937 Marijuana Act
 - * Around the same time as morphine and opioid development
- CBD and THC considered schedule I narcotic
 - * 2018 CBD not considered schedule I
- Come a long way with CBD
 - * NIH funds studies on CBD
 - * WHO: August 17, 2018 – no dependence, no public health problems
 - * FDA: May 2018: - no abuse potential

81

THC, CBD, Hemp – The Basics in more detail...

Cannabinoids: THC, CBD

- **THC** (delta-9-Tetrahydrocannabinol): psychoactive
 - Only compound in cannabis family that will get you “high”
- Main active compound in cannabis; will give positive drug test

82

Types of Cannabinoids

Endocannabinoids (EC): brain-derived

- Sources: omega-3, omega-6
- Anandamide (AEA)- “bliss chemical”

Phytocannabinoids: plant-derived

- Sources: buds, extracts, etc. of THC, **CBD**

Synthetic cannabinoids: lab-derived

- Examples: THC (Dronabinol/Marinol, Syndros, Cesamet); CBD (Cannabidiol/Epidiolex)

83

Synthetic cannabinoids: lab-derived

- Not great at mimicking nature
- Peer reviewed and systematic reviews concluded
 - Lower efficacy
 - Increased risk of adverse effects than phytocannabinoids
- Much higher affinity for CB1 and CB2 receptors than THC
 - Decrease therapeutic response
 - Decrease tolerability
 - Increased psychosis, paranoia, and side effects

84

So now that EVERYONE is selling it and talking about it...

- How do cannabinoids work?
 - Endocannabinoid (EC) system
 - CB1 and CB2 receptors that impact memory, pain, inflammation, appetite, immune system
 - CB1: CNS, genitourinary system, eyes, peripheral neurons, adrenals, heart, lung
 - CB2: CNS, immune system (spleen, tonsils, lymph nodes, thymus), bones, eyes, heart, gut

85

CB1 and CB2 Receptors

- THC – agonist to the CB1 and CB2 receptors and higher affinity
 - This is why THC comes with the risk of bad side effects
 - Anxiety, dysphoria, psychosis, sedation, subjective intoxication
 - THC can slow the development of frontal lobe with binding (agonist)
 - Not good for young brains, frontal lobe not developed until 21-25 years old
 - Nociceptive pain – mask the symptoms
- CBD – antagonist activity and lower affinity
 - Save for immature frontal lobe
 - No intoxication, euphoria, or paranoia (in normal doses)
 - Anti-inflammatory action
- THC and CBD do not cause respiratory depression or heart attack like opioid risks

86

3 Types of Products in the Market

- Full Spectrum CBD
 - Contains trace amounts of THC (delta 9 THC)
 - Should include other cannabinoid compounds
 - Multiple cannabinoids and terpenes
 - Lower dose than isolate by 5-10 times
 - Stable shelf life
 - Might fail a work or drug recovery program drug test – avoid
- Broad Spectrum CBD
 - No detectable THC
 - Other phytochemicals, terpenes
 - Won't fail a drug test
- Isolate CBD
 - Only CBD
 - Least medical benefits
 - Won't fail a drug test
 - Need high doses – 5-10 times more than full spectrum
 - Unstable shelf life

Doesn't work for everyone and everything
But CBD has a broad spectrum of uses

87

Starting to See Outcomes of Studies

Remember illegal until 2018

88

Fibromyalgia

- Allopathic way to treat is Cymbalta, Lyrica, and Savella
 - 8-10% say really effective
- Full spectrum cannabinoids – 62% very effective

89

Hemp Derived CBD Full Spectrum with Opioids

- 97 patients
- 15 mg softgels, average dose 30 mgs
- 53% of patients stopped or decreased opioid use in 8 weeks
- 94% reported better sleep or decrease pain
- CBD could significantly reduce opioid use and improve sleep quality

90

CBD with Drug Addiction

- Decreases reward facility effect and seeking behavior in opioid dependence
 - Not cocaine
- Decreases opioid seeking behavior
- Potential for relapse prevention in cocaine and alcohol

91

CBD to Recommend Need

- Dosing
- Delivery
- Interactions
- Monitoring
- Side effects
- Tolerability
- Risks
- Product selection

92

What to Look in a Company

- "Medical grade CBD"
- Certificate of Analysis (COA) – ask questions
 - Lot specific, comprehensive, is the lab iso-certified for cannabinoids
 - Checking heavy metals
 - Checking for molds, fungus, and bacteria
 - Manufacturing process
 - Planting process
 - Indoor or outdoor
 - Using pesticides
- The spectrums they have
 - If have isolate – does they do stability testing

93

Practical Application- Side Effects

- ⌚ Elevated LFTs (liver function tests)
 - * AST and ALT – why *THESE!!!*
 - * High doses of CBD
 - ⌚ Drowsiness/Dizziness
 - ⌚ Diarrhea
 - ⌚ Dry mouth
 - ⌚ Hypotension
 - ⌚ Increase in IOP
 - ⌚ Change in appetite
- Generally, side effects are most often seen in people taking HIGH doses of CBD

94

Our Associations Fought Hard

We took this course for a reason

95

Cases Where I Recently Used My DEA


96

Conditions Which May Require Pain Management


- ⌚ Large cornea abrasions
 - * Cornea burn
 - * PRK/PTK
- ⌚ Orbital trauma
- ⌚ Orbital blowout fractures
- ⌚ Scleritis




97



98



99



100



101

 <p>Optometric Education Consultants</p>	<p>Thank You!</p> <p>Treatment of Pain Opioid Choices and Issues for Patient and Practitioner</p> <p>Greg Caldwell, OD, FAAO Mid-Winter Getaway Scottsdale 2022 Sunday, February 2022 </p>
---	---

102