



Optometric
Education
Consultants



Neural Pearls

Joseph Sowka, OD, FAAO, Diplomate



1


Disclosure

Dr. Joseph Sowka has been a speaker for Bausch + Lomb and on the advisory board for Zeiss, and a co-owner of Optometric Education Consultants. He has no direct financial interest in any of the diseases, products or instrumentation mentioned in this presentation.

The ideas, concepts, conclusions and perspectives presented herein reflect the opinions of the speaker; he has not been paid, coerced, extorted or otherwise influenced by any third-party individual or entity to present information that conflicts with his professional viewpoints.

2

Know what can kill, maim, and blind immediately.




3




Once you **have** ruled out bad stuff, you have time to figure things out.

4

Acute, painful ANYTHING is a neuro-ophthalmic emergency.



5




Urgency of evaluation is dictated by duration of condition

6

46 YOM

- Reports waking up 3 months ago not being able to see OD
- LP OD, 20/20 OS
- Disc pallor OD- no other concurrent findings
- Last medical exam unknown- no medical hx
- Resident gets nervous- sends to ER immediately
- How long do we have to get this worked up?



7

What is the bad stuff?

- GCA
 - Any sudden vision loss in the elderly
- Pituitary apoplexy
 - Headache, field loss, diplopia
- Aneurysm
 - Pupils
- Papilledema
 - Clinical suspicion
- Carotid dissection
 - Horner syndrome

8

66 YOF**ESR = 96**

- New onset sudden vision loss
 - VA: 20/200 (longstanding macular scar)
 - Noticed inferior vision loss x 1 day
 - Inferior arcuate scotoma
- OD disc edema- mild pallor, no hemorrhages or telangiectasia
- OS disc- small, crowded disc at risk; C/D < 0.2
- Mild headache- relieved by OTC
- Malaise and loss of appetite- lost 7 lbs over 4 weeks
- No jaw claudication or temporal head pain
- What to do?

9




ANY acute vision loss in the elderly is GCA until proven otherwise

10

Anterior ISCHEMIC OPTIC NEUROPATHY

- Hypoperfusion of the posterior ciliary arterial supply to the anterior optic nerve head.
- May be arteritic (AAION) or non-arteritic (NAAION)
- Mechanical factors and atherosclerotic disease play a role in the non-arteritic form while vasculitis contributes in the arteritic form.
- Unilateral presentation but high incidence of subsequent contralateral involvement
 - AAION


11

AAION vs NAAION

12

NAAION

- Risk factors:
 - Hypertension, diabetes, atherosclerotic disease, small optic nerves
- Inferior field defects
- Hyperemic swollen nerve- disc at risk
- Progressive moderate vision loss with potential recovery
- Late 30s/ early 40s and beyond
- Painless




OptometricEdu.com/webinars

13

AAION

- Pallid optic nerve swelling with flame hemorrhages, arteriole attenuation and NFL infarcts
- Pain (of some sort)
- Severe optic nerve dysfunction
- Visual field defects
- Giant cell arteritis/ PMR- risk factors
- Typically 70s, uncommon under 60
 - Any patient over 50 is at risk
- High risk bilateral involvement



OptometricEdu.com/webinars

14

Diagnosis

- Careful history: Must directly ask about nonvisual symptoms
 - Headache (present in over 90%), scalp tenderness, jaw claudication (almost diagnostic), ear pain, arthralgias, temple pain and/or tenderness, malaise, intermittent fevers
- Examination
- Laboratory studies
 - Erythrocyte sedimentation rate
 - Lowered by statins and NSAIDs
 - C-reactive protein
 - Not affected by statins and NSAIDs
 - Elevated platelet count

OptometricEdu.com/webinars

15

Initial symptoms in GCA

A NORMAL EXAM

- Headache
- PMR
 - Chair and stair
- Fever
- Visual symptoms without vision loss
 - TIA, diplopia
- Weakness, malaise, fatigue
- What do these things have in common?

OptometricEdu.com/webinars

16

Vision Loss and Ocular Findings in GCA


- AION
- CRAO
- PION
- TIA
- Transient diplopia
- CWS

OptometricEdu.com/webinars

17

Headache and pain in GCA

- Temporal
- Occipital
- Neck
- Ear
- Jaw
- Scalp



OptometricEdu.com/webinars

18

AAION

Diagnosis

- Prodrome, GCA symptoms
- Elevated ESR/CRP (combination of the two gives high specificity [97%])
- Elevated platelet count (acute phase reactant)
- Ophthalmoscopy
- Fluorescein angiography
- Temporal artery Biopsy
 - Negative biopsy: Read the report: "No giant cells, no active arteritis"
 - Focal interruption of the internal elastic lamina => healed arteritis

Treatment

- Prompt steroids and hydration
- Consider IV when vision loss present
 - Very effective in prevention of second eye
 - Occasionally restores vision
 - Best done through ER
 - 250 mg solumedrol QID x 3 days followed by orals

OptometricEdu.com/Webinars

19

AAION versus NAAION

- Think AAION >> NAAION
 - Systemic symptoms of GCA
 - TVOs/TIAs
 - AION + cilioretinal artery occlusion
 - Evidence of posterior ciliary artery occlusion on FA
 - Multiple unexplained CWS
 - Early massive vision loss
 - Bilateral simultaneous vision loss
 - Chalky white optic disc edema
- Elevated
 - ESR/CRP
 - Platelets

OptometricEdu.com/Webinars

Hayreh SS, Podhajsky PA, Zimmerman B. Occult giant cell arteritis: ocular manifestations. *Am J Ophthalmol*. 1998 Apr;125(4):521-6.

20



NAAION is diagnosed in the negative. Get the tests done promptly with AION

21



Remember the E's in GCA:
Elderly, ESR is Elevated, only sees the Big E on the eye chart, and its an Emergency

22

Neuro-ophthalmic Urgencies and Emergencies

- GCA ✓
 - Any sudden vision loss in the elderly
- Pituitary apoplexy
 - Headache, field loss, diplopia
- Aneurysm
 - Pupils
- Papilledema
 - Clinical suspicion
- Carotid dissection
 - Horner syndrome

OptometricEdu.com/Webinars


23

28 YOF

- Presents with intermittent blurred vision & visual "grey-outs", intermittent horizontal diplopia, and chronic headache steadily worsening X 2 weeks
- MHx: "white coat hypertension", shoulder injury X 6 mos
- Meds: Flexeril" 10 mg BID PRN
- Height / weight: 5'3", 220 lbs.
- VA: OD 20/20, OS 20/20
- Pupils & motility: normal

OptometricEdu.com/Webinars

24



25

28 YOF

- Additional hx: Dull "ringing" in ears
- BP: 142/100
- SLE: unremarkable
- T₄: OU 16 mm Hg
- VF: blind spot enlargement & nasal step defect OU
- Serology Normal
- Imaging: small ventricles, otherwise normal
- LP: O.P. = 510 mm H₂O; all CSF studies normal
- DX: Pseudotumor cerebri (PTC)

OptometricEdu.com/webinars

27

PAPILLEDEMA: Signs & Symptoms


- **Signs:**
 - bilateral disc edema
 - superior & inferior aspects of discs affected FIRST
 - obliteration of optic cup
 - hemorrhages common
 - absence of SVP
 - Paton's folds
 - highly variable VF defects
 - enlarged blind spot (early)
 - arcuate defects and constricted (late)
 - NO RAPD typically
 - VA near normal
- **Symptoms:**
 - **Visual:**
 - transient visual obscurations
 - intermittent horizontal diplopia
 - **General:**
 - headache common
 - nausea & vomiting
 - dizziness
 - tinnitus

OptometricEdu.com/webinars

28

Papilledema Types:

- **Acute**
 - Hemorrhages, exudates, hyperemia, RNFL edema
- **Chronic**
 - Minimal hemorrhage/exudate. Collateral vessels may be present
- **Atrophic**
 - Eventually occurs if papilledema remains chronic. Optic disc pallor



OptometricEdu.com/webinars

29

PAPILLEDEMA Pathophysiology


- Disc edema results from axoplasmic stasis
 - intracellular fluids, metabolic by-products accumulate and are regurgitated at the level of the optic nerve head
 - in papilledema, cerebral edema is effectively transmitted along the common meningeal sheaths of the brain and optic nerve producing an engorged, swollen disc.

OptometricEdu.com/webinars

30

PAPILLEDEMA Pathophysiology

- Associated with intracranial abnormalities:
 - increased brain volume (intracranial mass lesion)
 - increased intracranial blood volume
 - intracranial hemorrhage
 - increased CSF volume
 - Hydrocephalus
 - Ventricular blockage by mass lesion



OptometricEdu.com/webinars

31

PAPILLEDEMA Management

- Rule out "swollen disc masqueraders"
 - ultrasonography can be invaluable in differentiating ONHD
 - also consider color, margins, SVP, vasculature, etc.
- Acute papilledema constitutes a medical emergency
 - Immediate neuro-imaging to rule out an intracranial mass.
 - If imaging is normal, lumbar puncture to measure CSF pressure and exclude meningitis or other disease processes is necessary.
- Atrophic papilledema with significant vision/field loss:
 - urgent measures must be undertaken to prevent blindness
- Papilledema accompanied by any neurologic abnormalities, fever or stiff neck:
 - Possible serious underlying neurologic abnormality, intracranial infection or bleed requiring immediate medical attention.

32

PTC vs. IIH

- Pseudotumor Cerebri (PTC)
 - Increased intracranial pressure in the absence of an intracranial mass lesion
 - Many causative agents have been identified
- Idiopathic Intracranial Hypertension (IIH)
 - Increased intracranial pressure without an identifiable cause
 - Young, obese females are at risk
- Primary PTC
 - IIH
- Poor CFS drainage

33

PSEUDOTUMOR CEREBRI DIAGNOSIS

- SI/SX: consistent with increased ICP
- Papilledema
- Normal neurological examination
 - except for cranial nerve 6 abnormalities
- Neuro-imaging: Normal without evidence of hydrocephalus, mass, or structural lesion, thrombosis
- Normal CSF composition
 - Elevated LP opening pressure
 - Adults: > 250 mm CSF
 - Children: > 280 mm CSF
 - > 250 mm CSF if not sedated/obese

34

PSEUDOTUMOR CEREBRI DIAGNOSIS

- LP may be deferred if:
 - MRI/MRV shows no additional abnormalities and has characteristic findings of flattened globe and empty sella.
- No evidence of fever or acute infection
- Typical profile

35

PSEUDOTUMOR CEREBRI MANAGEMENT


- No visual loss
 - Symptomatic headache therapy
 - Acetazolamide 500 mg tid
 - Weight reduction
- Mild visual loss
 - Acetazolamide 500 mg tid
 - Furosemide, Topiramate, Zonisamide
 - Weight reduction

36

PSEUDOTUMOR CEREBRI MANAGEMENT

- No/ Mild visual loss
 - Prognosis
 - Excellent (all signs and symptoms, visual loss)
 - 6-9 months
 - Follow-up and visual fields
- Role of weight loss
 - Treat the primary problem
 - 10% weight loss
 - Prevent recurrence
 - Keep the weight down

37




38

33 YOF


- Horizontal diplopia
- Headache
- TVOs 20/day
- Denies OCP, tetracyclines, vitamin A
- Lost 10 lbs- headaches improved
- 118/72
- 5'5"; 160lbs; BMI 26.62

OptometricEdu.com/webinars

39



40



41

PSEUDOTUMOR CEREBRI MANAGEMENT

- Severe, or progression of visual loss
 - Optic nerve sheath decompression (ONSD)
 - High-dose IV steroids and acetazolamide
 - Lumboperitoneal shunt
 - Failed ONSD
 - Declined ONSD
 - Intractable headache

OptometricEdu.com/webinars

42

IIH is a slowly progressive condition...until its not.

43

Fulminant IIH

- Same diagnostic criteria for IIH/ PTC
- Less than 4 weeks between symptoms and severe loss of acuity or field
- Vision rapidly worsening over several days
- Typically need CSF diversion surgery and/ or ONH sheath fenestration

OptometricEdu.com/webinars

44

Neuro-ophthalmic Urgencies and Emergencies

- GCA ✓
 - Any sudden vision loss in the elderly
- Pituitary apoplexy
 - Headache, field loss, diplopia
- Aneurysm
 - Pupils
- Papilledema ✓
 - Clinical suspicion
- Carotid dissection
 - Horner syndrome

OptometricEdu.com/webinars


45

63 YOIM


- Long standing glaucoma patient
- Sudden onset of orbital pain x 3 days
- + DM; +HTN
- On coumadin
- Pacemaker
- No vision change
- Presents as walk-in emergency glaucoma eval

OptometricEdu.com/webinars

46



47



48



5 mm
unresponsive

2 mm
responsive

49

63 YOIM

- Pupil involved CN III palsy
- 3 days duration at least
- Most likely cause: intracranial aneurysm
- Sent to ED with detailed notes and recommendations
- Endovascular therapy with coils
- Hospitalized 23 days

50



Secondary aberrant regeneration

51

CN III Palsy Clinical Picture

- An eye that is down and out with a ptosis
- Adduction, elevation, depression deficits
- Isocoric or anisocoric



52

83 YOM

- Diabetic: LBS in 300s; a1C around 11
- Pupils normal
- MRI ordered through PCP
- "Indications for imaging: Brain ischemia"
- What 2 mistakes are here?



53



The world's best neuroradiologist cannot help you if you don't order the scan, order the correct scan, and tell them what to look for

54

Neuroimaging for the primary care OD

- Disclosure: I do not read MRIs (There are ODs that do – I'm not one of them)
 - What you don't know can hurt you a whole lot
 - That's the reason for residencies in radiology and specialties in neuroradiology
 - Thinking that I am as good is irresponsible (e.g. neuroradiologist identifying ciliary body on MRI)
- Rules for ECP: order the correct scan and read the report to ensure that the right thing was done
 - If you have questions, doubts, or concerns, reach out to the radiologist
 - Form a relationship with a local imaging center- find out about the practice
 - Some have better results with MRA and others with CTA

55

What to order, how, and why



- Disc edema/ suspect papilledema: Brain MRI with and without contrast looking for mass lesion, hydrocephalus, hemorrhage, flattened globe, empty sella; MRV looking for cerebral venous sinus thrombosis.
- Optic nerve/chiasmal disease: MRI orbits and chiasm with and without contrast with fat suppression
 - Snowball in a snowstorm
- Optic neuritis/suspect MS: MRI orbits and chiasm with and without contrast with fat suppression; MRI brain with and without contrast.
- Horner Syndrome: Brain MRI with and without contrast; CTA (or MRA) head and neck looking for cerebral artery dissection; MRI chest with lung apex and brachial plexus
 - Horner protocol or sympathetic plexus
- Suspected aneurysm (CN 3 palsy): CTA/CT and MRA/MRI with concentration to Circle of Willis
 - If high risk aneurysm- send to ER and tell them what to do.
- Don't just send to the ER without helping them. They won't get it right.

OptometricEdu.com/webinars

56

CN III Anatomy

- Vulnerable to compression from aneurysm in subarachnoid space
 - Posterior communicating artery (PCOM)
 - Junction PCOM and ICA
 - Tip of basilar artery



OptometricEdu.com/webinars

57

Still More Clues



- Pupil involved CN III palsy is PCOM aneurysm until proven otherwise
- Incomplete palsy is PCOM aneurysm until proven otherwise
 - Regardless of pupil
- 30% of CN III palsy are caused by aneurysm
- Pain is pain
 - Only helpful when not present
- Vasculopathic CN III will resolve in time
- Life threatening posterior communicating aneurysm will rupture in time

OptometricEdu.com/webinars




58

Still More Clues

- CN III palsy caused by aneurysm
 - 20% die within 48 hrs from rupture
 - 50% overall die
 - Average time from onset to rupture – 29 days
 - 80% rupture w/i 29 days
 - Many never make it to hospital
- Ruptured aneurysms
 - 5% surgical mortality
 - 60% functional impairment post-op
- Unruptured aneurysms
 - No mortality; 75% with normal outcomes; 50% with CN III recovery

OptometricEdu.com/webinars

59



Never dilate a patient with a cranial nerve 3 palsy

60


61

Rules for CN III palsy imaging

- High suspicion of aneurysm: DSA (gold standard)
- CT/CTA is preferred non-invasive imaging for CN III palsy
 - CT for SAH
- CTA requires contrast- renal impairment prefers MRI/MRA
- CTA superior to MRI when patient can't have MRI
 - Pacemaker, claustrophobia
- MRI superior for non-aneurysmal causes (tumor)
 - MRA adds very little time to scan

OptometricEdu.com/webinars

62



63


A Different patient and Prognosis

- 63 YOF
- Diabetes and HTN
- Sudden onset retro-orbital pain



OptometricEdu.com/webinars

64



65

Which is better? One or two?



Resolves over several weeks



Hospitalized 23 days with 2 neurosurgical procedures

66

Suspect the worst

- Optometrist sees patient with CN III palsy
- Referred to ophthalmologist next day
- Pt dies from SAH before consult

OptometricEdu.com/webinars

67

Does presence of vasculopathic risk factors help?

- Arteriosclerotic risk factors in elderly favors microvascular etiology but does not rule out aneurysm
- HTN, DM, atherosclerosis, hypercholesterol all common and don't protect against aneurysm
- Answer: **no**, but makes me very nervous when NOT present

OptometricEdu.com/webinars

68

Does acuteness of presentation help?

- Ans: **Yes and No**
- Aneurysm expansion usually produces acute manifestations, but chronic and evolving cases well known
- Acute is more worrisome
- Chronic and improving less worrisome but does not rule out aneurysm
- Resolved without recurrence reassuring

OptometricEdu.com/webinars

69

Aneurysm Risk Assessment: Isolated CN 3 palsy

- | | |
|------------------------------|-----------|
| • Isolated dilated pupil | none |
| • Complete CN3-normal pupil | low |
| • Partial CN3 – normal pupil | high |
| • Pupil involved CN3 | emergency |

OptometricEdu.com/webinars

70

Never out of the woods

- Pt develops CN III palsy from aneurysm
- Successfully treated with aneurysm clip
 - All coils are inert and MRI safe; not all clips are MRI safe
- Radiologic tech doesn't verify type of clip
- Pt undergoes F/U MRI with non-MRI safe clip in major medical center
- Clip displaces during MRI
- Patient has fatal hemorrhage during procedure
- Patient survived disease...killed by follow up

OptometricEdu.com/webinars

71

Neuro-ophthalmic Urgencies and Emergencies

- | | |
|---|----------------------|
| • GCA ✓ | • Aneurysm ✓ |
| • Any sudden vision loss in the elderly | • Pupils |
| • Pituitary apoplexy | • Papilledema ✓ |
| • Headache, field loss, diplopia | • Clinical suspicion |
| | • Carotid dissection |
| | • Horner syndrome |

OptometricEdu.com/webinars

72

39 YOM

- Previous history of migraine- developed a new and different headache.
- He presented to a hospital emergency room where he underwent a non-contrast enhanced computed tomography (CT) and magnetic resonance imaging (MRI) which were subsequently interpreted as normal.
 - His headache was attributed to migraine, and he was medicated as such and discharged.
- Three days later, he developed horizontal and vertical diplopia and presented urgently.

OptometricEdu.com/webinars

73



74

39 YOM

- His visual acuity and visual fields were normal.
- He manifested a right pupil-sparing, external partial cranial nerve three palsy and concurrent right sixth nerve palsy. He also complained of worsening headache and lethargy.
- Where is the lesion?
- Let's contact the radiologist for a second reading...

OptometricEdu.com/webinars

75

39 YOM

- He was immediately sent for repeat imaging to include contrast-enhanced MRI of the parasellar area and MRA to rule out intracavernous aneurysm and pituitary apoplexy.
- Imaging revealed a pituitary macroadenoma with intratumor hemorrhage consistent with pituitary apoplexy.
- Lateral spread into the right cavernous sinus and possible spread into the left cavernous sinus as well.
- No mass effect on the optic chiasm or prechiasmatic intracranial portion of the optic nerve.
 - Hence normal acuity and fields
- The patient was immediately admitted for endocrinological and neurosurgical evaluation

OptometricEdu.com/webinars

76

Pituitary apoplexy

- Pituitary apoplexy is a severe and potentially fatal medical condition complicating 2-12% of pituitary adenomas and characterized by the variable association of headache, vomiting, visual impairment, ophthalmoplegia, altered mental state and consciousness, lethargy, and panhypopituitarism.
- Hemodynamic instability may be result from adrenocorticotrophic hormone deficiency, which can be fatal.
- Occurs due to a rapid expansion, mainly caused by hemorrhage or infarction of a preexisting (known or unknown) adenoma

OptometricEdu.com/webinars

77

Pituitary apoplexy

- Most common presenting symptom occurring in 90 % of patients is sudden onset of severe headache
 - Commonly described as frontal or retro-orbital.
 - Pituitary apoplexy is often overlooked as a possible cause of "thunderclap headache" where diagnostic evaluations tend to direct to more common causes of this presentation including subarachnoid hemorrhage, cerebral venous sinus thrombosis, and cervical artery dissection.
- Approximately 50% have visual abnormalities
 - Blurred vision
 - Cranial nerve palsy (CN III) or palsies
 - Cranial nerve VI most common, followed by CN III
 - Visual field defects
 - Bitemporal hemianopsia
 - Facial weakness

OptometricEdu.com/webinars

78

Pituitary apoplexy

- Most symptomatic patients undergo CT scanning in an emergency setting due to the clinical suspicion of acute intracranial hemorrhage
- Acute hemorrhagic infarct may be seen on CT
 - Non-hemorrhagic infarcts will usually show no abnormalities without intravenous contrast
- MRI with contrast is the most effective imaging in cases of suspected pituitary apoplexy
 - MRI is superior to CT

OptometricEdu.com/webinars

79

Pituitary apoplexy

- Positive outcome in most cases
- Conservative medical treatment
 - Stabilize and replace diminished pituitary hormones
- Surgical decompression
 - Trans-sphenoidal or subfrontal transcranial approach
 - Patients with visual impairment and neuro-ophthalmic dysfunction will be selected for surgery.
- Patient was medically stabilized, and surgery delayed due to COVID lock down
- Ultimately underwent successful surgical decompression

OptometricEdu.com/webinars

80

Neuro-ophthalmic Urgencies and Emergencies

- GCA ✓
 - Any sudden vision loss in the elderly
- Pituitary apoplexy ✓
 - Headache, field loss, diplopia
- Aneurysm ✓
 - Pupils
- Papilledema ✓
 - Clinical suspicion
- Carotid dissection
 - Horner syndrome

OptometricEdu.com/webinars


81

33 YOWM


- Occipital HA x 4 mos
 - Visual aura with HA
- Worsens when standing after sitting
- Relieved by sleep
- Denies vision loss, nausea, diplopia, pain on eye movement, behavioral changes
- 20/20 OD, OS with myopic correction
- Pupils, EOMs, conf fields normal OU
- Biomicroscopy normal OU
- IOP 12 mm Hg OU
- Nasally obliquely inserted nerves

OptometricEdu.com/webinars

82



83



84

33 YOWM

- Co-manage with PCP- internist
- MRI w and w/o contrast of brain and orbits
- Complete blood work blood work up including FTA-ABS/RPR ; Lyme titer; CBC w/differential
- Rule out mass lesion, infections, collagen vascular and autoimmune etiology.

OptometricEdu.com/webinars

85

33 YOWM

- MRI
- Pt had MRI done and mass was identified in fronto/parietal region more toward right side
- Outcome?

86



A normal visual field does not mean that there isn't anything wrong

87

Case

- A 25-year-old woman was involved in a minor automobile accident where she was hit by another driver. The accident was reportedly minor, with no initial injury to either driver, and both cars were able to be driven away. She felt that she experienced only a mild-to-moderate bump during the accident with no head trauma or loss of consciousness. However, immediately upon waking the next morning, though she had no physical pain, she experienced profound double vision.

88

What is the likely cause?

- A subarachnoid hemorrhage
- A third nerve palsy
- Orbital fracture
- Fourth nerve palsy
- Sixth nerve palsy

89

Case

- She described the diplopia as vertical and worse at near. She had a distinct right hyper deviation which, on alternate cover test, worsened in left gaze and right head tilt. This was a signature motility of a cranial IV (trochlear) palsy.




90



91

CN IV Anatomy

- Exits the midbrain posteriorly and decussates
- Longest course
- Travels around tentorium, through cavernous sinus, through SOF
- Most prone to trauma




OptometricEdu.com/webinars

92

CN IV Palsy

- Longstanding CN IV palsy may present with diplopia from decompensation
 - Observe old photos for head tilt (*Facebook Tomography*)
- Rule of 40-30-20-10
 - 40% traumatic
 - 30% idiopathic
 - 20% vascular
 - 10% CNS lesion



OptometricEdu.com/webinars

93

CN IV Management

- Isolated, non-traumatic:
 - Evaluate for ischemic diseases
 - Non-ischemic causes of non-traumatic, isolated CN IV palsy rare
- Look for longstanding decompensation
 - Increased vertical vergences
 - Old photos

OptometricEdu.com/webinars

94

73 YOM

- New onset vertical diplopia
- Left 4th nerve palsy
- Relieved by 2 PD BD
- *"Doc, I also noticed that my gripper is off"*
 - Mild left-handed weakness
- Medical history: Treated lung cancer
 - Currently on maintenance chemotherapy
- Approach and outcome?

OptometricEdu.com/webinars

95



Never diagnose idiopathic (or ischemic) anything in a patient with a history of cancer

96

77 YOM

- Pt and sister presents insistent on cataract surgery.
 - BVA 20/40 OD, 20/70 OS with commensurate NS
- Chronic horizontal diplopia that has recently gotten worse (2 weeks)
- Left abduction deficit ~ 40%
- Medical history: Inoperable chondrosarcoma with lysis of clivus extending to left petrous apex and occipital condyle with sphenoidal, ethmoidal and temporal bone involvement. Compression of jugular vein. Has undergone ~ 50 radiation treatments.
- Vocal paralysis
- Cranial nerve IX, X, XII palsies
- Imaging obtained: CT with contrast (pacemaker precludes MRI)- new soft mass in left nasopharynx
 - Likely squamous cell carcinoma

OptometricEdu.com/webinars

97

CN VI Palsy

- Hallmark sign is horizontal diplopia, greater at distance, with an abduction deficit



OptometricEdu.com/webinars

98

CN VI Palsy

- Check motilities at distance
- Forced duction testing




OptometricEdu.com/webinars

99

More About Mass Lesions

- CN VI is stretched against the clivus
- CN VI palsy common in ICP rises/mass lesions/ PTC
- Bilateral CN VI palsy and disc edema is indicative of mass lesions and increased intracranial pressure



OptometricEdu.com/webinars

100

CN VI Management

- Each case of CN VI palsy should be classified as traumatic or non-traumatic.
- Non-traumatic cases should be subdivided as neurologically isolated (just CN VI palsy) or non-neurologically isolated (something else).
- Additionally, patients should be ascribed to one of 3 groups: children, young adults, and older adults

OptometricEdu.com/webinars


101

CN VI Demographic Groups

- Older adults (**usually not bad**)
 - Vascular disease common- resolves-3mos
 - Consider GCA over 60 yrs
- Children (**may be bad**)
 - Presumed viral illness, trauma, malignancy (50%)
- Young adults (**usually bad**)
 - Vascular disease (4%) and idiopathic (13%) uncommon
 - Usually complicated CN VI palsy (hemiparesis, Horner syndrome, facial paresis)
 - Cerebrovascular accidents involving the pons, aneurysm (typically within the cavernous sinus) or neoplasm (33%-cavernous sinus, pons), **MS (24%)**.

OptometricEdu.com/webinars

102



103

CN VI Palsy in Older Adults

- In cases of isolated CN VI palsy in older adults with a history of diabetes or hypertension, neuroimaging and other extensive evaluation can be deferred, unless the palsy progresses, fails to improve over 3 months, or other neurologic complications develop.
- Ischemic vascular palsies typically progress over several days, but progression over two weeks warrants neuroimaging.

OptometricEdu.com/webinars

104



When looking for mass cause of CN VI palsy, the base of the pons and cavernous sinus are two common hiding spots

105



If you are watching a presumptive ischemic CN VI palsy and you are wrong, you likely have not hurt the patient.

106

78 YOF

- Sudden onset of ptosis OS
- Immediately following parathyroid surgery
- Headache and eye pain
- Dilation lag and positive Iopidine test
- This is when things get complicated...



OptometricEdu.com/webinars

107

Discussion

- What is Horner's Syndrome?
 - a triad of clinical signs arising from disruption of sympathetic innervation to the eye and ipsilateral face that causes *miosis*, upper lid *ptosis*, mild elevation of the lower lid, and *anhidrosis* of the facial skin.

108

What is the most likely cause?

- Lung cancer
- Carotid dissection
- Direct surgical trauma to the nerve
- Migraine

OptometricEdu.com/webinars

109

Horner's Syndrome: Etiologies

- First-order neuron disorder: Stroke (e.g., verteobasilar artery insufficiency or infarct); tumor; multiple sclerosis (MS), and, rarely, severe osteoarthritis of the neck with bony spurs.
- Second-order neuron disorder: Tumor (e.g., lung carcinoma, metastasis, thyroid adenoma, neurofibroma). Patients with pain in the arm or scapular region should be suspected of having a Pancoast tumor. In children, consider neuroblastoma, lymphoma, or metastasis.



110

Horner's Syndrome: Etiologies


- Third-order neuron disorder: Headache syndrome (e.g., cluster, migraine, Raeder paratrigeminal syndrome), internal carotid dissection, herpes zoster virus, otitis media, Tolosa-Hunt syndrome, neck trauma/tumor/inflammation, prolactinoma.
- Other rare causes: Cervical paraganglioma, ectopic cervical thymus



111

Carotid Dissection

- A 3rd-order Horner's and ipsilateral head, eye, or neck pain of acute onset should be considered diagnostic of internal carotid dissection unless proven otherwise.



OptometricEdu.com/webinars

112

Carotid Dissection

- Carotid artery dissection presents with the sudden or gradual onset of ipsilateral neck or hemicranial pain, including eye or face pain
- Often associated with other neurologic findings including an ipsilateral Horner's syndrome, TIA, stroke, anterior ischemic optic neuropathy, subarachnoid hemorrhage, or lower cranial nerve palsies
 - 52% with ocular or hemispheric stroke with 6 days
 - 67% within first week; 89% within 2 weeks; none after 31 days
- Horner's from suspected carotid dissection should go to ER

OptometricEdu.com/webinars

113



If you listen to the patient, they will tell you the diagnosis

114

Case History 73 YOWF

- CC: swollen left eyelid x 3 months
- Not happy with previous doctor
 - "They aren't listening to me"
- Highly allergic person- had pain and ear blockage on right side of face while gardening- thinks something got into her eye
- Rx Zylet, Azasite, oral antihistamines, hot and cold compresses- no improvement
- PCP tested for GCA- negative
- Presumed allergic reaction
 - No itching, persistent and unilateral
- Hypothyroid, smoker

115



116



119

Neuro-ophthalmic Urgencies and Emergencies

- GCA ✓
 - Any sudden vision loss in the elderly
- Pituitary apoplexy ✓
 - Headache, field loss, diplopia
- Aneurysm ✓
 - Pupils
- Papilledema ✓
 - Clinical suspicion
- Carotid dissection ✓
 - Horner syndrome

OptometricEdu.com/Webinars

120

78 YOM

- Acute onset diplopia, blurred vision, and dilated pupils
- Went to ED- worked up for stroke
- CT/CTA MRI/MRI all normal
 - Reviewed reports- everything in order
- Referred by colleague after exam
- Vision improving, pupils less dilated, endpoint nystagmus, non-specific horizontal diplopia
- Pt on anti-muscarinic for bladder
- Mydriasis and blurred vision = ??
- What about that history?
- Outcome?



OptometricEdu.com/Webinars

121

Take Care with VESicare

- 56 YOF
- Treated glaucoma
- IOP suddenly in mid 30s with meds
- Bilateral blurred vision and dilated pupils

OptometricEdu.com/Webinars

122

The Case of the Curbside Consult

- Conference focusing on neurological issues
- 48 YOF- wife of society president
- Complaints of memory loss and loss of smell and taste (Pre-COVID)
 - Fearful of neurodegenerative condition.
- Hx of tonsillectomy two months earlier
- Facial and persistent jaw pain
- Seen by multiple specialists (neuro, ENT, PCP)
 - Put on multiple medications including at least 2 oral antibiotics
 - Personal experience with Mucinex
- Outcome?

OptometricEdu.com/Webinars

124



Its not always a brain tumor. Think about medication toxicity

125

Toxicities

- Blurred vision and dilated pupils
- Ethambutol
 - Toxic optic neuropathy
- Amiodarone/ Pacerone
 - Toxic optic neuropathy
- Vigabatrin
 - Anticonvulsant for refractory focal epilepsy in children 2 years of age or older
 - May cause permanent, concentric peripheral visual field loss, thought to be secondary to drug-induced injury to both the retinal photoreceptors and the retinal ganglion cells and their axons.

 OptometricEdu.com/webinars

126

Immediately referring to the Emergency Department is acceptable management...if you are willing to help




127

Best handled in the ED*

- Suspected GCA
- Suspected aneurysm
- Suspected papilledema
- Suspected pituitary apoplexy
- Suspected carotid dissection
- CRAO/ BRAO/ TIA

- *As long as you are willing to help

 OptometricEdu.com/webinars

128