





1




2




3



4



7



8

### Pregnancy and Lactation Labeling Rule-FDA December 4, 2014, Final Rule

- Effective June 30, 2015
  - ★ Effective now for new medications and a 3-5-year phase in period (application)
- Labeling for human prescription drugs and biological products will include:
  - ★ Pregnancy
  - ★ Lactation
  - ★ Females and Males of Reproductive Potential
- Pregnancy (8.1)
  - ★ Pregnancy Exposure Registry – omit if not applicable
  - ★ Risk Summary – required subheading
    - Dose adjustments during pregnancy and the postpartum period - omit if not applicable
    - Maternal adverse reactions - omit if not applicable
    - Fetal/Neonatal adverse reactions - omit if not applicable
    - Labor or delivery - omit if not applicable
  - ★ Data - omit if none of the headings are applicable
    - Human Data - omit if not applicable
    - Animal Data - omit if not applicable

9

### Pregnancy and Lactation Labeling Rule-FDA December 4, 2014, Final Rule

- Lactation (8.2)
  - ★ Risk Summary - required subheading
  - ★ Clinical Considerations - omit if not applicable
  - ★ Data - omit if not applicable
- Females and Males of Reproductive Potential (8.3) - omit if none of the headings are applicable
  - ★ Pregnancy testing - omit if not applicable
  - ★ Contraception - omit if not applicable
  - ★ Infertility - omit if not applicable

10

### Pre-June 30, 2015

11

### Post-June 30, 2015

12

### Post-June 30, 2015

13


### Dacryocystitis

Test me your answer 814-931-2030  
Patient as a severe allergic reaction to Penicillin and Keflex (Epipen)  
Which antibiotic would you use?

- A. Augmentin
- B. Azithromycin
- C. Cephalexin
- D. Bactrim
- E. Cipro


14

## Antibiotic Paradigm Thru 2019




15

## Antibiotic Paradigm Changed 2020



16


## Cross Reaction



17

## "Sulfa Antibiotic Allergy"

- ~ If a true allergy: Mast cell and IgE mediated, causing hives, itch, congestion, lip and tongue swelling, tightness in throat, and bronchial constriction
- \* Anaphylaxis
- ~ The "S" is typically not the hapten/antigen
- ~ The arylamine group is the hapten/antigen




18

## What to ask about allergies

- ~ For the patient
  - \* When did you react?
  - \* What was the reaction?
    - Systemic versus topical

- ~ For you...
  - \* For sulfa allergic patients
    - Antibiotic structure sulfa allergy
      - Sulfamethoxazole/trimethoprim
    - Non-antibiotic structure agents
      - Carbonic anhydrase inhibitors
      - Hydrochlorothiazide
      - Celecoxib (Celebrex)
      - Glipizide, glyburide, glimepiride



19

## Amoxicillin + Clavulanic acid (Augmentin)

- ~ "Uber Amoxicillin"
- ~ Kills everything, good for everyone
  - 12 weeks old and older
- ~ Safe in pregnancy...category B
- ~ Watch for PCN allergies
- ~ Adults: 250, 500 and 875 mg
  - \* Dose of clavulanate varies
- ~ Children <100 pounds: oral suspension 25-45 mg/kg divided into 2 doses
- ~ Covers *Staph*, *Strep* and *Haemophilus influenzae*

20

### Azithromycin (Zithromax)

- ~ Macrolide antibiotic (erythromycin is prototype)
- ~ Drug of choice in PCN sensitive patients
- ~ All age groups and pregnancy category B
- ~ No renal adjustment
- ~ Adult:
  - \* 250 mg bid (day1), 250 mg qd (day 2-5), 6 pack
  - \* 500 mg qd x 3 days, tri-pack
- ~ Children <16: 10 mg/kg (day1), 5 mg/kg (day 2-5)
- ~ Covers *Staph*, *Strep* and *Haemophilus influenzae*
- ~ Better tolerated than erythromycin, little GI upset
- ~ Chlamydia...1 g qd

21

### Zithromax (azithromycin)

- ~ "The Vegas Drug"- Chlamydia...1 g qd



22

### Cephalexin (Keflex)


- ~ Cross reaction with PCN sensitive patients
  - \* Approximately 3 - 10%
- ~ 1st generation, moderately effective against PCN-ase
- ~ Good for Gram +, +/- for *Haemophilus* (-)
- ~ Available in 250 and 500 mg
- ~ Category B
- ~ Adult: typically, 500 mg bid x 1 week
  - \* Maximum 4 g in 24 hrs
- ~ FYI...Drug of choice for blow out fractures

23

### Cefuroxime (Ceftin)

- ~ 2nd generation
- ~ Better for *Haemophilus* (-)
- ~ Children: 3 months to 12 years old, oral suspension 15 mg/kg divided into 2 doses x 10 days
- ~ Available in 125, 250 and 500 mg
  - \* FYI: adults typically 250 mg bid x 10 days
- ~ Category B

24



### Cross Reaction

25

### Sulfa Drugs

- ~ Limited use...last line of defense
- ~ Bactrim SS
  - \* 400 mg sulfamethoxazole/ 80 mg trimethoprim
  - \* 1-2 tab PO bid
- ~ Bactrim DS (double strength)
  - \* 800 mg sulfamethoxazole/ 160 mg trimethoprim
  - \* 1 tab PO bid

26



### Ciprofloxacin (Cipro) Levofloxacin (Levaquin)


- ~ In my opinion, an end of the line, antibiotic to use...allergic to PCN, cephalosporins, macrolides...
- ~ Really effective, because they are BROAD
- ~ Would avoid if pregnant, BF, and in kids
  - \* Only use 18 years or older (oral)
- ~ Cipro and Levaquin available in 250, 500 and 750 mg
  - \* Cipro 750 mg for only severe infections (usually life-threatening pneumonia)
- ~ 500 mg bid x 1 week-Cipro
- ~ 500 mg qd x 1 week-Levaquin
- ~ Levaquin-tendon ruptures

27

### Fluoroquinolone ADRs

- ~ Retinal detachment (1 per 2,500 pts)
  - \* WHAT???
  - Mechanism is possible through destruction of collagen and connective tissue...
- ~ QT prolongation in newer agents
- ~ Photosensitivity
- ~ Tendon rupture
  - \* Watch shoulders, wrists, Achilles


28




### Dacryocystitis

Text me your answer 814-931-2030  
Patient as a severe allergic reaction to Penicillin and Keflex (Epipen)  
Which antibiotic would you use?

- Augmentin
- Azithromycin
- Cephalexin
- Bactrim
- Cipro



29



~ Remember...patient allergic to PCN and Keflex

### Treatment

- \* Polytrim QID OD
- \* Zithromax
  - Disp: 5 day z-pak
  - Use as directed PO

Text me your answer 814-931-2030

- ~ Dilation and Irrigation
  - \* Contraindication or indication?
- ~ Confirmed nasolacrimal duct blockage
  - \* DCR, dacryocystorhinostomy

30

### What group of antibiotics are we missing?

31

Text me your answer 814-931-2030

When we use doxycycline or minocycline for dry eye disease, ocular surface, and/or meibomian gland dysfunction. What property of this pharmaceutical are we benefitting from?

- Bacteriostatic action – anti-infective
- Anti-protozoan action – Malaria
- Works for Methicillin-resistant Staphylococcus aureus (MRSA)
- Anti-inflammatory action – inhibiting bacterial lipases
- I don't know

32

## Treatment Failure

- ~ If you continue to think of doxycycline and minocycline as antibiotics, treatment failure will be the result
- ~ From this point on consider them a steroid

33

48-year-old man  
OU red, gritty, sandy and dry feeling

Va 20 / 20  
20  
cc 20

Current Correction  
R -2.00 sphere  
L -3.00 sphere

EOMS: full, unrestricted PERRL (-)APD  
CF: ortho D/N CF: full by FC OU

34



- ~ Diagnosis
  - \* Rosacea
- ~ What findings support your diagnosis?
  - \* Telangiectasias
  - \* Erythema of the cheeks, forehead and nose
  - \* Rhinophyma
    - Indicates chronic
- ~ Let us get a closer look

35

## A Closer Look




36

## Rosacea Blepharitis (Inflammatory Blepharitis, MGD)

- ~ Diagnosis?
- ~ Treatment?
  - \* In my opinion, most under treated condition
  - \* Warm compresses
  - \* Lid hygiene
  - \* Artificial tears
  - \* Omega 3 fatty acid
    - EPA and DHA total 1500 mg (1000 mg minimum)
  - \* Dermatological consult (Acne Rosacea)
  - \* Oral antibiotics...???
  - Which one and why??

37

## Tetracycline Analog



38

## Minocycline / Doxycycline

- ~ Drug of choice for marginal inflammatory blepharitis, posterior blepharitis, MGD, evaporative disease
- ~ AB, anti-inflammatory and anti-collagenase
- ~ No renal adjustment
- ~ 50-100 mg qd-bid 2-12 weeks (pulse)
  - \* Lower maintenance dose
- ~ 20 mg Periostat (Doxycycline)
  - \* Helpful in those with stomach or GI sensitivity
  - \* Excellent for those requiring long maintenance dose

39

## My Paradigm for Minocycline / Doxycycline

- |   |   |
|---|---|
| <ul style="list-style-type: none"> <li>~ Status of MG                             <ul style="list-style-type: none"> <li>* Insipissated</li> <li>* Turbid</li> <li>* Clear</li> </ul> </li> </ul> | <ul style="list-style-type: none"> <li>~ Minocycline / Doxycycline Paradigm                             <ul style="list-style-type: none"> <li>* Maximum dosage for 2-12 weeks (pulse)                                     <ul style="list-style-type: none"> <li>□ 100 mg BID, QD</li> </ul> </li> <li>* 50-100mg qd while turbid</li> <li>* 20 mg longer treatments                                     <ul style="list-style-type: none"> <li>□ Periostat (Doxycycline)</li> </ul> </li> <li>* 20 mg if maintenance dose needed</li> </ul> </li> </ul> |
|---|---|


40

## Customize Treatment

- ~ 50 mg Minocycline with pill cutter (25 mg)
- ~ Oracea- 40 mg of Doxycycline total
  - \* 30 mg immediate release
  - \* 10 mg sustained release
- ~ AzaSite (azithromycin ophthalmic solution) 1.0%
  - \* Initiate early in treatment
  - \* Adjunctive when patient is already on Doxycycline
  - \* Alternative in states that do not have oral therapeutic licensure

41

## Successfully Treated



- ~ Warm Compresses
- ~ Lid Scrubs
- ~ Artificial Tears, Systane Balance
- ~ Omega 3 (1500 EPA and DHA)
- ~ Mino 100 mg PO 6 weeks, 50 mg 3 months, 20 mg maintenance (Doxycy)
- ~ Steroids, Tobradex qid (5 weeks with taper)
  - \* Moderately red and thickened lid margins
  - \* Marginal infiltrates

42

## Hyclate vs Monohydrate

- ~ I get calls from the pharmacist
  - \* Doxycycline
    - Doryx
      - Enteric coated hyclate pellet
    - Adoxa
      - Monohydrate



43

## Doxycycline and Minocycline Adverse Drug Reactions

- ~ Enhanced photosensitivity
- ~ Avoid in children and pregnancy (Category D)
- ~ Can interfere with how penicillin kills bacteria
- ~ Enhances the effects of
  - \* Coumadin
  - \* Digoxin
- ~ Idiosyncratic Secondary intracranial hypertension
  - \* Pseudotumor cerebri
  - \* Isotretinoin (Accutane) - for severe acne and doxy/mino increase this risk
- ~ Hyperpigmentation
  - \* Antacids that contain aluminum, calcium, magnesium, bismuth subsalicylate, and iron may inhibit doxy/mino
  - \* Same for seizure meds like barbiturates, carbamazepine, phenytoin



44


Text me your answer 814-931-2030

I have seen a patient with optic nerve head edema from doxycycline or minocycline:


- A. Yes
- B. No
- C. I don't know

45

Benign intracranial hypertension  
"It's not rare if it's in your chair"



46




47

Minocycline Optic Nerve Edema



48

Minocycline Optic Nerve Edema



49

OMG



50





51



52

### Delafloxacin™ (Baxdela)

- ~ A fluoroquinolone antibiotic for acute bacterial skin and skin structure infections
- ~ Approved 2017 – Melinta Therapeutics
- ~ Available orally and intravenously
- ~ Adverse effects: SO NEW...but in clinical trials, the only ophthalmic side effects that were noted = blurred vision!
  - Only time will tell if retinal detachment is something to worry about with this new FQ!

53

### Nuzyra™ (omadacycline)

- ~ Approved 2018
- ~ Tetracycline antibiotic
- ~ Approved for PO/IV treatment of patients with bacterial skin infections or community-acquired bacterial pneumonia
- ~ Chelation issues JUST like other tetracyclines!
- ~ ADRs: Nausea, vomiting, diarrhea, constipation, insomnia

54

### Seysara™ (sarecycline)

- ~ Approved 2018
- ~ Tetracycline drug
- ~ Indicated for the treatment of inflammatory acne in non-nodular, moderate to severe acne vulgaris
- ~ Can be taken WITH or WITHOUT food!
- ~ ADRs: nausea

55

### Xerava™ (eravacycline)

- ~ Approved 2018
- ~ Tetracycline antibiotic
- ~ Indicated for the treatment of intra-abdominal infections in adults
- ~ IV ONLY

56

## Orals in Herpetic Eye Disease

- ~ Valtrex
- ~ Acyclovir
- ~ Famvir
- ~ Neurontin
- ~ Lyrica
- ~ Doxycycline
- ~ L-Lysine
- ~ Tagamet
- ~ Tricyclic antidepressants
  - Amitriptyline, nortriptyline

57

## Fun Facts About Herpes

- ~ Are a leading cause of human viral disease
  - Second only to influenza and cold viruses
- ~ There are more than 130 herpes viruses identified
  - 8 infect humans (9 if you count HHV-6A and HHV-6B as two separate)
  - 5 infect the eye
    - Herpes simplex 1
    - Herpes simplex 2
    - Varicella zoster
    - Epstein Barr
    - Cytomegalovirus
- ~ USA 25% of the population is seropositive for HSV by 4 years old
  - Nearly 100% are seropositive by age 60
  - Lifetime prevalence of ocular manifestation in all HSV infected people is 1%

58

## 8 Humans- 5 Eye

Viruses of Humans	Common Name	Subfamily	Viruses of Humans	Common Name	Subfamily
Human herpesvirus 1	Herpes simplex type 1	alpha	Human herpesvirus 1	Herpes simplex type 1	alpha
Human herpesvirus 2	Herpes simplex type 2	alpha	Human herpesvirus 2	Herpes simplex type 2	alpha
Human herpesvirus 3	Varicella zoster	alpha	Human herpesvirus 3	Varicella zoster	alpha
Human herpesvirus 4	Epstein Barr	gamma	Human herpesvirus 4	Epstein Barr	gamma
Human herpesvirus 5	Cytomegalovirus	beta	Human herpesvirus 5	Cytomegalovirus	beta
Human herpesvirus 6/7	roseola infantum	beta	Human herpesvirus 6/7	roseola infantum	beta
Human herpesvirus 8	Kaposi's Sarcoma agent	gamma	Human herpesvirus 8	Kaposi's Sarcoma agent	gamma

59

## Herpes Viruses are Classified by Their Location in the Latent State

Human Herpes Virus Type	Name	Sub Family	Target Cell Type	Latency	Transmission
1	Herpes simplex 1 (HSV-1)	Alphaherpesvirinae	Neurons	Neurons	Oral contact
2	Herpes simplex 2 (HSV-2)	Alphaherpesvirinae	Neurons	Neurons	Oral contact, sexually transmitted
3	Varicella Zoster Virus (VZV)	Alphaherpesvirinae	Neurons	Neurons	Contact or respiratory route
4	Epstein-Barr Virus (EBV)	Gammaposvirinae	B lymphocytes, epithelial	B lymphocytes	Saliva
5	Cytomegalovirus (CMV)	Cytomegalovirinae	Epithelial, monocytes, lymphocytes	Monocytes, lymphocytes, and possibly others	Contact, nasal, breastfeeding, transplacental, congenital
6	Human herpesvirus 6 (HHV-6)	Betaherpesvirinae	T lymphocytes and others	T lymphocytes and others	Contact, respiratory route
7	Human herpesvirus 7 (HHV-7)	Betaherpesvirinae	T lymphocytes and others	T lymphocytes and others	Unknown
8	Kaposi's Sarcoma-associated herpesvirus (KSHV)	Gammaposvirinae	Endothelial cells	Unknown	Exchange of body fluids?

60

## Herpes Simplex Virus Keratitis

- ~ Is a leading cause of corneal blindness in the United States
  - Primarily caused by HSV-1 (65%)
- ~ Keratitis nomenclature
  - ~ Infectious epithelial keratitis
    - It's not critical to determine HSV 1 or 2
  - ~ Stromal keratitis
  - ~ Endothellitis
  - ~ Neurotrophic keratopathy

61

## Varicella-Zoster Virus (VZV)

- ~ AKA: Herpes Zoster Virus or Herpes Human Virus 3
- ~ Vesicles on tip of nose indicate nasociliary involvement
  - High risk of ocular manifestations



62

### Varicella-Zoster Virus (VZV)

~ The best time to diagnose and treat



63



64

### Text me your answer 814-931-2030

Ever wonder why a Primary Care Physician sends you with Herpes Zoster already on oral Valtrex and prednisone?

- A. Yes
- B. No
- C. Never happened to me

65

### Varicella-Zoster Virus (VZV)



- ~ Vesicles on tip of nose indicate nasociliary nerve involvement
  - High risk of ocular manifestations
- ~ Ocular findings associated with VZV
  - \* Episkleritis
  - \* Scleritis
  - \* Keratitis
  - \* Uveitis
  - \* Iris atrophy
  - \* Glaucoma
  - \* Vitritis
  - \* Retinitis
  - \* Choroiditis
  - \* Optic neuritis
  - \* CN palsy

66

### Renal Impairment

- ~ Identify patients on hemodialysis
- ~ Adjustment made by patient's creatinine clearance (CrCl)...ml/min
  - \* Work with patient's PCP/Internist

67

### Zovirax (acyclovir)

- ~ Good for simplex and zoster
- ~ Available in 200, 400 and 800 mg, IV
- ~ Dosage: 800 mg/5 times/day (4 grams daily)
  - ~ Poor GI absorption
- ~ Maintenance dose: 200-400 mg bid
- ~ Caution if impaired renal function
  - \* Excreted by kidneys
- ~ Category B

68

## Off-Label

- ✓ Valtrex and Famvir used for the eye
- ★ Off label
- ★ Only approved for genital herpes
- ★ Won't find dosage in PDR for ocular usage

69

## Famvir (famciclovir)

- ✓ Available in 125, 250 and 500 mg
- ✓ Dosage: Zoster 500 mg tid
- Recurrent Simplex 125-250 mg bid
- ✓ Caution if impaired renal function
- ✓ Category B
- ✓ No longer available via Novartis in USA as brand name

70

## Valtrex (valacyclovir)

- ✓ Pro-drug of acyclovir
- ✓ Available in 500 and 1000 mg
- ✓ GI upset
- ✓ HSV-1, HSV-2, VZV
- ✓ Dosage: 1g tid x 1 week (3 grams daily)
- ✓ Caution if impaired renal function
- ✓ Category B

71

## Beside the dosing frequencies...

- ✓ What is different about the oral antivirals?



- ✓ Main reason for early discontinuation of oral acyclovir in HEDS
- ✓ Gastrointestinal side effects
- ✓ Rash

Many patients on oral acyclovir have GI symptoms

72

## Text me your answer 814-931-2030

- Which oral anti-viral is lactose free?
- Acyclovir
  - Famvir
  - Valtrex
  - They all have lactose

73

## Acyclovir vs. Valacyclovir vs. Famciclovir

### What is the difference?

ACYCLOVIR is the first name for acyclovir, a synthetic nucleoside analogue active against Herpesvirus, HSV-1, HSV-2, VZV, and Epstein-Barr virus. It is used for the treatment of genital herpes, shingles, and chickenpox. Each capsule contains 200 mg of acyclovir and the active ingredients are acyclovir, hydroxyethylamine, and sodium lauryl sulfate. The capsule shell contains gelatin, PEG 400, and titanium dioxide. It is not recommended for use in patients with known hypersensitivity to acyclovir or any of the other ingredients.

VALACYCLOVIR (valacyclovir hydrochloride) is the hydrochloride salt of the L-valeryl ester of acyclovir. It is a synthetic nucleoside analogue active against Herpesvirus, HSV-1, HSV-2, VZV, and Epstein-Barr virus. It is used for the treatment of genital herpes, shingles, and chickenpox. Each capsule contains 500 mg of valacyclovir and the active ingredients are valacyclovir, hydroxyethylamine, and sodium lauryl sulfate. The capsule shell contains gelatin, PEG 400, and titanium dioxide. It is not recommended for use in patients with known hypersensitivity to valacyclovir or any of the other ingredients.

FAMCICLOVIR (famciclovir hydrochloride) is the hydrochloride salt of the L-famciclovir ester of acyclovir. It is a synthetic nucleoside analogue active against Herpesvirus, HSV-1, HSV-2, VZV, and Epstein-Barr virus. It is used for the treatment of genital herpes, shingles, and chickenpox. Each capsule contains 250 mg of famciclovir and the active ingredients are famciclovir, hydroxyethylamine, and sodium lauryl sulfate. The capsule shell contains gelatin, PEG 400, and titanium dioxide. It is not recommended for use in patients with known hypersensitivity to famciclovir or any of the other ingredients.

Generics available in the US contain lactose

\* In Europe you can get generic famciclovir without lactose (Teva Pharmaceuticals, Israel)

Zovirax® contains lactose  
Presence or absence of lactose in generic acyclovir tablets

Valtrex® and all generics are free of lactose

74



Acyclovir vs. Valacyclovir vs. Famciclovir  
What is the difference?


CNS Effects in Elderly Patients

- Acyclovir and valacyclovir carry a higher risk of CNS adverse effects in the elderly:
  - Agitation
  - Hallucinations
  - Confusion
- Clinical Take Home Point:
  - Consider famciclovir in older patients who CNS side effects with acyclovir or valacyclovir
  - Other major concern with elderly patients is age-related reduced kidney function

75

24-48 hours


- Zirgan
- Viroptic
- Orals only
- Orals and Amniotic Membrane




76

Is there a difference in efficacy between topical and orals in the various forms of ocular herpes?

Ganciclovir ophthalmic gel



Oral antivirals:  
Acyclovir  
Valacyclovir  
Famciclovir



77

Epithelial keratitis  
There seems to be equivalence


Oral acyclovir (Zovirax) in herpes simplex dendritic corneal ulceration

60 patients with HSV dendritic ulceration included a small number with stromal involvement keratitis randomized to oral vs. topical acyclovir

No statistically significant difference in time to resolution (mean = 5 days)

"Oral acyclovir alone appeared as effective as topical antiviral therapy in the treatment of simplex epithelial keratitis."

Oral delivery appears to get to corneal target even though it is an avascular tissue!




78

Vaccines

- Zostavax™ – live vaccine; 60 years and older
  - "the only game in town..."
  - 50-60% effective; 1 dose
  - Efficacy wanes after 4-5 years
- Shingrix™ – has replaced Zostavax™
  - We are moving in the right direction!
  - Recommended for 50 years and older
  - 90+% effective; 2 doses; IM; recombinant vaccine
  - Efficacy seems solid up to 7-8 years

79




Optometric  
Education  
Consultants

Oral Pharmaceuticals in Eye Care  
"Pearls from an Optometrist and Pharmacist"

Greg Caldwell, OD, FAAO  
Tracy Offerdahl, PharmD, Bpharm, RPh, FAAO

CE Sarasota 2023  
Optometric Education Consultants  
Saturday, March 4, 2023



80