

NEURAL PEARLS

Joseph Sowka, OD, FAAO, Diplomate

DISCLOSURE:

Dr. Joseph Sowka is/ has been in the past 24 months a consultant or member of the advisory or speaker boards for B&L. All relevant relationships have been mitigated. Dr. Sowka has no direct financial interest in any of the diseases, products or instrumentation mentioned in this presentation.

He is a co-owner of Optometric Education Consultants.
www.optometricedu.com

DO NOT BE AN ACADEMIC TERMINATOR

He can't be bargained with...he can't be reasoned with... he doesn't care what time it is... he doesn't care about the break...he doesn't care whose time he runs over... he will absolutely not stop until his message is delivered

NEURO-OP IS A FINANCIALLY REWARDING SPECIALTY...

NEURO-OP IS A FINANCIALLY REWARDING SPECIALTY...SAID NOBODY EVER

Specialty	Number of Claims
Cardiology	250
General	220
Neurology	220
Orthopedics	100
Urology	80
Neuro	40
Endocrinology	30
Other	20

Specialty	Average Indemnity (\$)
Endocrinology	100
Orthopedics	25
Genes	5
Neurology	1
Neuro	1
Urology	10
General	50
Reflexive	4
General	40
Reflex	30

Courtesy of Ophthalmic Mutual Insurance Company



Neuro-Op

- High risk
- Is this urgent? Can it wait?
- Complicated
- Diagnose and Adios
- Schedule-busting



KNOW WHAT CAN KILL, MAIM, AND BLIND
IMMEDIATELY.



ONCE YOU HAVE RULED OUT BAD STUFF, YOU
HAVE TIME TO FIGURE THINGS OUT.



ACUTE PAINFUL IS A NEURO-
OPHTHALMIC EMERGENCY



ACUTE PAINFUL ANYTHING IS A NEURO-
OPHTHALMIC EMERGENCY






US EYE

URGENCY OF EVALUATION IS DICTATED BY DURATION OF CONDITION

US EYE

46 YOM

- Reports waking up 3 months ago not being able to see OD
- LP OD, 20/20 OS
- Disc pallor OD- no other concurrent findings
- Last medical exam unknown- no medical hx
- Resident gets nervous- sends to ER immediately
- How long do we have to get this worked up?



US EYE

ER PHYSICIAN ENCOUNTERING A PATIENT WITH AN EYE PROBLEM



US EYE

Neuro-ophthalmic Urgencies and Emergencies

- GCA
 - Any sudden vision loss in the elderly
- Pituitary apoplexy
 - Headache, field loss, diplopia
- Aneurysm
 - Pupils
- Papilledema
 - Clinical suspicion
- Carotid dissection
 - Horner's syndrome

US EYE

66 YOF

ESR = 96

- New onset sudden vision loss
 - VA: 20/400 (longstanding macular scar)
 - Noticed inferior vision loss x 1 day
 - Inferior arcuate scotoma
- OD disc edema- mild pallor, no hemorrhages or telangiectasia
- OS disc- small, crowded disc at risk; C/D < 0.2
- Mild headache- relieved by OTC
- Malaise and loss of appetite- lost 7 lbs over 4 weeks
- No jaw claudication or temporal head pain
- What to do?

US EYE




ANY ACUTE VISION LOSS IN THE ELDERLY IS GCA UNTIL PROVEN OTHERWISE



Anterior ISCHEMIC OPTIC NEUROPATHY


- Hypoperfusion of the posterior ciliary arterial supply to the anterior optic nerve head.
- May be arteritic (AAION) or non-arteritic (NAAION)
- Mechanical factors and atherosclerotic disease play a role in the non-arteritic form while vasculitis contributes in the arteritic form.
- Unilateral presentation but high incidence of subsequent contralateral involvement
 - AAION

AAION VS NAAION




NAAION

- Risk factors:
 - Hypertension, diabetes, atherosclerotic disease, small optic nerves
- Inferior field defects
- Hyperemic swollen nerve- disc at risk
- Progressive moderate vision loss with potential recovery
- Late 30s/ early 40s and beyond
- Painless



AAION

- Pallid optic nerve swelling with flame hemorrhages, arteriole attenuation and NFL infarcts
- Pain (of some sort)
- Severe optic nerve dysfunction
- Visual field defects
- Giant cell arteritis/ PMR- risk factors
- Typically 70s, uncommon under 60
 - Any patient over 50 is at risk
- High risk bilateral involvement



Diagnosis

- Careful history: Must directly ask about nonvisual symptoms
 - Headache (present in over 90%), scalp tenderness, jaw claudication (almost diagnostic), ear pain, arthralgias, temple pain and/or tenderness, malaise, intermittent fevers
- Examination
- Laboratory studies
 - Erythrocyte sedimentation rate
 - Lowered by statins and NSAIDs
 - C-reactive protein
 - Not affected by statins and NSAIDs
 - Elevated platelet count

Initial symptoms in GCA

- Headache
- PMR
 - Chair, hair, stair, fair
- Fever
- Visual symptoms without vision loss
 - TIA, diplopia
- Weakness, malaise, fatigue
- What do all of these things have in common?


A normal exam

Vision Loss and Ocular Findings in GCA

- AION
- CRAO
- PION
- TIA
- Transient diplopia

Headache and pain in GCA

- Temporal
- Occipital
- Neck
- Ear
- Jaw
- Scalp



AAION

Diagnosis

- Prodrome, GCA symptoms
- Elevated ESR/CRP (combination of the two gives high specificity [97%])
- Elevated platelet count (acute phase reactant)
- Ophthalmoscopy
- Fluorescein angiography
- Temporal artery Biopsy
 - Negative biopsy: Read the report: "No giant cells, no active arteritis"
 - Focal interruption of the internal elastic lamina= healed arteritis
- Temporal artery ultrasound

Treatment

- Prompt steroids and hydration
- Consider IV when vision loss present
 - Very effective in prevention of second eye
 - Occasionally restores vision
 - Best done through ER
 - 250 mg solumedrol QID x 3 days followed by orals



NAAION IS DIAGNOSED IN THE NEGATIVE. GET THE TESTS DONE PROMPTLY WITH AION



**REMEMBER THE E'S IN GCA:
ELDERLY, ESR IS ELEVATED, ONLY SEES THE BIG E
ON THE EYE CHART, AND ITS AN EMERGENCY**

What to say to the ER doc

- Don't say, "This patient has blurred vision"




- Say, "This elderly patient has suddenly lost vision in one eye and will go totally blind from giant cell arteritis if they aren't treated with steroids immediately!"

Neuro-ophthalmic Urgencies and Emergencies

- **GCA**
 - Any sudden vision loss in the elderly
- Pituitary apoplexy
 - Headache, field loss, diplopia
- Aneurysm
 - Pupils
- Papilledema
 - Clinical suspicion
- Carotid dissection
 - Horner's syndrome

28 YOF

- Presents with intermittent blurred vision & visual "grey-outs", intermittent horizontal diplopia, and chronic headache steadily worsening X 2 weeks
- MHx: "white coat hypertension", shoulder injury X 6 mos
- Meds: Flexeril® 10 mg BID PRN
- Height / weight: 5'3", 220 lbs.
- VA: OD 20/20, OS 20/20
- Pupils & motility: normal



What questions do you want to ask?
What tests do you want to perform?

28 YOF


- Additional hx: Dull "ringing" in ears
- BP: 142/100
- SLE: unremarkable
- T₄: OU 16 mm Hg
- VF: blind spot enlargement & nasal step defect OU
- Serology Normal
- Imaging: small ventricles, otherwise normal
- LP: O.P. = 510 mm H₂O; all CSF studies normal
- DX: Pseudotumor cerebri (PTC)

PAPILLEDEMA: Signs & Symptoms

- Signs:
 - bilateral disc edema
 - superior & inferior aspects of discs affected FIRST
 - obliteration of optic cup
 - hemorrhages common
 - absence of SVP
 - Paton's folds
 - highly variable VF defects
 - enlarged blind spot (early)
 - arcuate defects and constricted (late)
 - NO RAPD typically
 - VA near normal
- Symptoms:
 - Visual:
 - transient visual obscurations
 - intermittent horizontal diplopia
 - General:
 - headache common
 - nausea & vomiting
 - dizziness
 - tinnitus

Papilledema Types:

- **Acute**
 - Hemorrhages, exudates, hyperemia, RNFL edema
- **Chronic**
 - Minimal hemorrhage/exudate. Collateral vessels may be present
- **Atrophic**
 - Eventually occurs if papilledema remains chronic. Optic disc pallor




PAPILLEDEMA Pathophysiology

- Disc edema results from axoplasmic stasis
 - Intracellular fluids, metabolic by-products accumulate and are regurgitated at the level of the optic nerve head
 - In papilledema, cerebral edema is effectively transmitted along the common meningeal sheaths of the brain and optic nerve producing an engorged, swollen disc.

PAPILLEDEMA Pathophysiology

- Associated with intracranial abnormalities:
 - Increased brain volume (intracranial mass lesion)
 - Increased intracranial blood volume
 - Intracranial hemorrhage
 - Increased CSF volume
 - Hydrocephalus
 - Ventricular blockage by mass lesion



PAPILLEDEMA Management

- Rule out "swollen disc masqueraders"
 - ultrasonography can be invaluable in differentiating ONHD
 - also consider color, margins, SVP, vasculature, etc.
- Acute papilledema constitutes a medical emergency
 - Immediate neuro-imaging to rule out an intracranial mass.
 - If imaging is normal, lumbar puncture to measure CSF pressure and exclude meningitis or other disease processes is necessary.
- Atrophic papilledema with significant vision/field loss:
 - urgent measures must be undertaken to prevent blindness
- Papilledema accompanied by any neurologic abnormalities, fever or stiff neck:
 - Possible serious underlying neurologic abnormality, intracranial infection or bleed requiring immediate medical attention.

PTC vs. IIH


- Pseudotumor Cerebri (PTC)
 - Increased intracranial pressure in the absence of an intracranial mass lesion
 - Many causative agents have been identified
- Idiopathic Intracranial Hypertension (IIH)
 - Increased intracranial pressure without an identifiable cause
 - Young, obese females are at risk
- Primary PTC
 - IIH
- Poor CSF drainage

PSEUDOTUMOR CEREBRI DIAGNOSIS

- SI/SX: consistent with increased ICP
- Papilledema
- Normal neurological examination
 - except for cranial nerve 6 abnormalities
- Neuro-imaging: Normal without evidence of hydrocephalus, mass, or structural lesion, thrombosis
- Normal CSF composition
 - Elevated LP opening pressure
 - Adults: > 250 mm CSF
 - Children: > 280 mm CSF
 - > 250 mm CSF if not sedated/obese

PSEUDOTUMOR CEREBRI DIAGNOSIS

- LP may be deferred if:
 - MRI/MRV shows no additional abnormalities and has characteristic findings of flattened globe and empty sella.
- No evidence of fever or acute infection
- Typical profile





PSEUDOTUMOR CEREBRI MANAGEMENT

- No visual loss
 - Symptomatic headache therapy
 - Acetazolamide 500 mg tid
 - Weight reduction
- Mild visual loss
 - Acetazolamide 500 mg tid
 - Furosemide, Topiramate, Zonisamide
 - Weight reduction

PSEUDOTUMOR CEREBRI MANAGEMENT



- No/Mild visual loss
 - Prognosis
 - Excellent (all signs and symptoms, visual loss)
 - 6-9 months
 - Follow-up and visual fields
- Role of weight loss
 - Treat the primary problem
 - 10% weight loss
 - Prevent recurrence
 - Keep the weight down

17 YOF; 5'4"; 155 lbs.; BMI 26.60; HA with exercise; 20/20 OD, OS 



33 YOF


- Horizontal diplopia
- Headache
- TVOs 20/day
- Denies OCP, tetracyclines, vitamin A
- Lost 10 lbs- headaches improved
- 118/72
- 5'5"; 160lbs; BMI 26.62



PSEUDOTUMOR CEREBRI MANAGEMENT



- Severe, or progression of visual loss
 - Optic nerve sheath decompression (ONSD)
 - High-dose IV steroids and acetazolamide
 - Lumboperitoneal shunt
 - Failed ONSD
 - Declined ONSD
 - Intractable headache



IIH IS A SLOWLY PROGRESSIVE
CONDITION...UNTIL ITS NOT

Fulminant IIH



- Same diagnostic criteria for IIH/PTC
- Less than 4 weeks between symptoms and loss of field/ acuity
- Vision worsening rapidly over several days
- Typically needs CSF diversion surgery and/or ONS fenestration

Neuro-ophthalmic Urgencies and Emergencies






- GCA
 - Any sudden vision loss in the elderly
- Pituitary apoplexy
 - Headache, field loss, diplopia
- Aneurysm
 - Pupils
- Papilledema
 - Clinical suspicion
- Carotid dissection
 - Horner syndrome

63 YOIM




- Long standing glaucoma patient
- Sudden onset of orbital pain x 3 days
- + DM; +HTN
- On coumadin
- Pacemaker
- No vision change
- Presents as walk-in emergency glaucoma eval





DX: Right pupil involved CN 3 palsy from aneurysm

- 50% die from aneurysm rupture w/i 29 days
- 20% die within 48 hours
- Needs emergency care and time counts



- Sent to ER with detailed notes, recommendations, and cell phone#. Called triage nurse in advance
- Pupil involved right third nerve palsy
- Most likely cause: intracranial aneurysm of posterior communicating artery
- Needs CT/CTA/ neurosurgical consult STAT
- Was in scanner within 45 minutes
- Leaking but unruptured aneurysm confirmed with CTA
- Endovascular therapy with coils successful (2 procedures)
- Hospitalized 23 days
- Ptosis improved, motility and pupil didn't, but he *did* live




CN III Palsy Clinical Picture

- An eye that is down and out with a ptosis
- Adduction, elevation, depression deficits
- Isocoric or anisocoric





83 YOM

- Diabetic; LBS in 300s;
- A1C around 11
- Pupils normal MRI ordered through PCP
- Indication for imaging: Brain ischemia
- What 2 errors were made here?



**THE WORLD'S BEST NEURORADIOLOGIST CAN'T
HELP YOU IF YOU DON'T ORDER THE SCAN,
ORDER THE RIGHT SCAN, AND TELL THEM WHAT
TO LOOK FOR.**



Still More Clues





EYE


- Pupil involved CN III palsy is PCOM aneurysm until proven otherwise
- Incomplete palsy is PCOM aneurysm until proven otherwise
 - Regardless of pupil
- **30% of CN III palsy are caused by aneurysm**
- Pain is pain
 - Only helpful when not present
- Vasculopathic CN III will resolve in time
- Life threatening posterior communicating aneurysm will rupture in time

Still More Clues

- CN III palsy caused by aneurysm
 - 20% die within 48 hrs from rupture
 - 50% overall die
 - Average time from onset to rupture – 29 days
 - 80% rupture w/ 29 days
 - Many never make it to hospital
- Ruptured aneurysms
 - 5% surgical mortality
 - 60% functional impairment post-op
- Unruptured aneurysms
 - No mortality; 75% with normal outcomes; 50% with CN III recovery




**NEVER DILATE A PATIENT WITH CRANIAL NERVE III
PALSY**



Rules for CN III palsy imaging

- High suspicion of aneurysm: DSA (gold standard)
- CT/CTA is preferred non-invasive imaging for CN III palsy
 - CT for SAH
- CTA requires contrast- renal impairment prefers MRI/MRA
- CTA superior to MRI when patient can't have MRI
 - Pacemaker, claustrophobia
- MRI superior for non-aneurysmal causes (tumor)
 - MRA adds very little time to scan
- Recent study shows majority of CN 3 palsy patients do not get the appropriate urgent imaging.



US EYE

A Different patient and Prognosis

- 63 YOF
- Diabetes and HTN
- Sudden onset retro-orbital pain

US EYE

Complete CN III palsy with pupil sparing and vasculogenic risk factors

WHICH IS BETTER? ONE OR TWO? US EYE

Resolves over several weeks

Hospitalized 23 days with 2 neurosurgical procedures

US EYE

Suspect the worst

- Optometrist sees patient with CN III palsy
- Referred to ophthalmologist next day
- Pt dies from SAH before consult

US EYE

Does presence of vasculopathic risk factors help?

- Arteriosclerotic risk factors in elderly favors microvascular etiology but does not rule out aneurysm
- HTN, DM, atherosclerosis, hypercholesterol all common and don't protect against aneurysm
- Answer: **no**, but makes me very nervous when NOT present

Does acuteness of presentation help?

- Ans: **Yes and No**
- Aneurysm expansion usually produces acute manifestations, but chronic and evolving cases well known
- Acute is more worrisome
- Chronic and improving less worrisome but does not rule out aneurysm
- Resolved without recurrence reassuring

Aneurysm Risk Assessment: Isolated CN 3 palsy

- Isolated dilated pupil none
- Complete CN3-normal pupil low
- Partial CN3 – normal pupil high
- Pupil involved CN3 **emergency**

What to say to the ER doc

- Don't say, "This patient has double vision"
- Say, "This patient has an aneurysm of the posterior communicating artery and is going to DIE if he doesn't get to neurosurgery immediately!"



Neuroimaging for the primary care OD

- Disclosure: I do not read MRIs (There are ODs that do- I'm not one of them)
 - What you don't know can hurt you a whole lot
 - That's the reason for residencies in radiology and subspecialties in neuroradiology
 - Thinking that I am as good is irresponsible (e.g. neuroradiologist identifying ciliary body on MRI)
- Rules for ECP: order the correct scan and read the report to ensure that the right thing was done
- If you have questions, doubts, or concerns, reach out to the radiologist
- Form a relationship with an imaging center- find out about the practice
 - Some have better results with MRA and others with CTA

What to order, how, and why

- Disc edema/suspect papilledema: Brain MRI with and without contrast looking for mass lesion, hydrocephalus, hemorrhage, flattened globe, empty sella; MRV looking for cerebral venous sinus thrombosis.
- Optic nerve/chiasmal disease: MRI orbits and chiasm with and without contrast with fat suppression
- Snowball in a snowstorm
- Optic neuritis/suspect MS: MRI orbits and chiasm with and without contrast with fat suppression; MRI brain with and without contrast.
- Homer Syndrome: Brain MRI with and without contrast; CTA (or MRA) head and neck looking for cerebral artery dissection; MRI chest with lung apex and brachial plexus
 - Homer protocol or sympathetic plexus
- Suspected aneurysm (CN 3 palsy): CTA/CT and MRA/MRI with concentration to Circle of Willis
 - If high risk aneurysm- send to ER and tell them what to do.
- Don't just send to the ER without helping them. They won't get it right.

Neuro-ophthalmic Urgencies and Emergencies

- **GCA**
 - Any sudden vision loss in the elderly
- Pituitary apoplexy
 - Headache, field loss, diplopia
- **Aneurysm**
 - Pupils
- **Papilledema**
 - Clinical suspicion
- Carotid dissection
 - Horner syndrome



39 YOM

- Previous history of migraine developed a new and worsening headache.
- He presented to a hospital emergency room where he underwent a non-contrast enhanced computed tomography (CT) and magnetic resonance imaging (MRI) which were subsequently interpreted as normal.
 - His headache was attributed to migraine, and he was medicated as such and discharged.
- Three days later, he developed horizontal and vertical diplopia



39 YOM

- His visual acuity and visual fields were normal.
- He manifested a right pupil-sparing, external partial cranial nerve three palsy and concurrent right sixth nerve palsy. He also complained of worsening headache and lethargy.
- Where is the lesion?
- Let's contact the radiologist for a second reading...



39 YOM

- He was immediately sent for repeat imaging to include contrast-enhanced MRI of the parasellar area and MRA to rule out intracavernous aneurysm and pituitary apoplexy.
- Imaging revealed a pituitary macroadenoma with intratumor hemorrhage consistent with pituitary apoplexy.
- Lateral spread into the right cavernous sinus and possible spread into the left cavernous sinus as well.
- No mass effect on the optic chiasm or prechiasmatic intracranial portion of the optic nerve.
 - Hence normal acuity and fields
- The patient was immediately admitted for endocrinological and neurosurgical evaluation



Pituitary apoplexy

- Pituitary apoplexy is a severe and potentially fatal medical condition complicating 2-12% of pituitary adenomas and characterized by the variable association of headache, vomiting, visual impairment, ophthalmoplegia, altered mental state and consciousness, lethargy, and panhypopituitarism.
- Hemodynamic instability may be result from adrenocorticotrophic hormone deficiency, which can be fatal.
- Occurs due to a rapid expansion, mainly caused by hemorrhage or infarction of a preexisting (known or unknown) adenoma



Pituitary apoplexy

- Most common presenting symptom occurring in 90 % of patients is sudden onset of severe headache
 - Commonly described as frontal or retro-orbital.
 - Pituitary apoplexy is often overlooked as a possible cause of "thunderclap headache" where diagnostic evaluations tend to direct to more common causes of this presentation including subarachnoid hemorrhage, cerebral venous sinus thrombosis, and cervical artery dissection.
- Approximately 50% have visual abnormalities
 - Blurred vision
- Cranial nerve palsy (CN III) or palsies
 - Cranial nerve VI most common, followed by CN III
- Visual field defects
 - Bitemporal hemianopsia
- Facial weakness

Pituitary apoplexy

- Most symptomatic patients undergo CT scanning in an emergency setting due to the clinical suspicion of acute intracranial hemorrhage
- Acute hemorrhagic infarct may be seen on CT
 - Non-hemorrhagic infarcts will usually show no abnormalities without intravenous contrast
- MRI with contrast is the most effective imaging in cases of suspected pituitary apoplexy
 - MRI is superior to CT

Pituitary apoplexy


- Positive outcome in most cases
 - Conservative medical treatment
 - Stabilize and replace diminished pituitary hormones
- Surgical decompression
 - Trans-sphenoidal or subfrontal transcranial approach
 - Patients with visual impairment and neuro-ophthalmic dysfunction will be selected for surgery.
- Patient was medically stabilized, and surgery delayed due to COVID lockdown
- Ultimately underwent successful surgical decompression

Neuro-ophthalmic Urgencies and Emergencies

- GCA
 - Any sudden vision loss in the elderly
- Pituitary apoplexy
 - Headache, field loss, diplopia
- Aneurysm
 - Pupils
- Papilledema
 - Clinical suspicion
- Carotid dissection
 - Horner syndrome


78 YOF

- Sudden onset of ptosis OS
- Immediately following parathyroid surgery
- Headache and eye pain
- Dilation lag and positive Iopidine test



WHAT IS HORNER'S SYNDROME? US EYE

A triad of clinical signs arising from disruption of sympathetic innervation to the eye and ipsilateral face that causes *miosis*, upper lid *ptosis*, mild elevation of the lower lid, and *anhidrosis* of the facial skin.



What is the most likely cause?

- Lung cancer
- Carotid dissection
- Direct surgical trauma to the nerve
- Migraine

HORNER'S SYNDROME: ETIOLOGIES



- First-order neuron disorder: Stroke (e.g., vertebralbasilar artery insufficiency or infarct); tumor; multiple sclerosis (MS), and, rarely, severe osteoarthritis of the neck with bony spurs.
- Second-order neuron disorder: Tumor (e.g., lung carcinoma, metastasis, thyroid adenoma, neurofibroma). Patients with pain in the arm or scapular region should be suspected of having a Pancoast tumor. In children, consider neuroblastoma, lymphoma, or metastasis.



HORNER'S SYNDROME: ETIOLOGIES




- Third-order neuron disorder: Headache syndrome (e.g., cluster, migraine, Raeder paratrigeminal syndrome), internal carotid dissection, herpes zoster virus, otitis media, Tolosa-Hunt syndrome, neck trauma/tumor/inflammation, prolactinoma.
- Congenital Horner's syndrome: Trauma (e.g., during delivery).
 - Heterochromia
- Other rare causes: Cervical paraganglioma, ectopic cervical thymus



Carotid Dissection



- A 3rd-order Horner's and ipsilateral head, eye, or neck pain of acute onset should be considered diagnostic of internal carotid dissection unless proven otherwise.



Carotid Dissection



- Carotid artery dissection presents with the sudden or gradual onset of ipsilateral neck or hemicranial pain, including eye or face pain
- Often associated with other neurologic findings including an ipsilateral Horner's syndrome, TIA, stroke, anterior ischemic optic neuropathy, subarachnoid hemorrhage, or lower cranial nerve palsies
 - 52% with ocular or hemispheric stroke with 6 days
 - 67% within first week; 89% within 2 weeks; none after 31 days
- Horner's from suspected carotid dissection should go to ER

What to say to the ER doc



- Don't say, "This patient has a little ptosis and a little pupil."



- Say, "This patient has a carotid artery dissection and will stroke out unless they get a CTA and referred to a stroke neurologist now!"

Neuro-ophthalmic Urgencies and Emergencies



- GCA
 - Any sudden vision loss in the elderly
- Pituitary apoplexy
 - Headache, field loss, diplopia
- Aneurysm
 - Pupils
- Papilledema
 - Clinical suspicion
- Carotid dissection
 - Horner syndrome



IF YOU LISTEN TO PATIENTS, THEY WILL TELL
YOU THE DIAGNOSIS

73 YOWF

- CC: swollen left eyelid x 3 months
- Not happy with previous doctor
 - “They aren’t listening to me”
- Highly allergic person- had pain and ear blockage on right side of face while gardening- thinks something got into her eye
- Rx Zylet, Azasite, oral antihistamines, hot and cold compresses- no improvement
- PCP tested for GCA- negative
- Presumed allergic reaction
 - No itching, persistent and unilateral
- Hypothyroid, smoker



Pre-tapidine

Post-tapidine



IMMEDIATELY REFERRING TO THE EMERGENCY
DEPARTMENT IS ACCEPTABLE MANAGEMENT...IF
YOU
ARE WILLING TO HELP

Best handled in the ED*

- Suspected GCA
- Suspected aneurysm
- Suspected papilledema
- Suspected pituitary apoplexy
- Suspected carotid dissection
- CRAO/ BRAO/ TIA



*As long as you are willing to help

