



Optometric
Education
Consultants

Treatment of Pain Opioid Choices and Considerations

Greg Caldwell, OD, FAAO

Mid-Winter Getaway
Optometric Education Consultants

Sunday, January 28, 2024



Disclosures- Greg Caldwell, OD, FAAO

All relevant relationships have been mitigated

- **Lectured for: Alcon, B&L, BioTissue, Dompé**
 - Disclosure: Receive speaker honorariums
- **Advisory Board: Dompé, ImmunoGen, Iveric**
 - Disclosure: Receive participant honorariums
- **I have no direct financial or proprietary interest in any companies, products or services mentioned in this presentation**
 - Disclosure: Non-salaried financial affiliation with Pharmanex
- **Healthcare Registries – Chairman of Advisory Council for Diabetes and AMD**
- **The content of this activity was prepared independently by me - Dr. Caldwell**
- **The content and format of this course is presented without commercial bias and does not claim superiority of any commercial product or service**
- **Optometric Education Consultants – Scottsdale, AZ, Pittsburgh, PA, Sarasota, FL , Barcelona, Spain, Orlando, FL, Mackinac Island, MI, Quebec City, Canada, and Nashville, TN- Owner**



My Practice

I am a clinician first then a scientist

- Some are scientists first then clinician
- I need to simplify for patient and patient care.
- Science is great, but not good if there isn't a clinical application.
- Some lectures are science based without clinical application.
- My lecture will be a hybrid. Showing clinical applications of the science



It is wonderful to have someone who's juggling so many aspects of optometry [scientific, clinical experience, teacher & lecturer]. It is refreshing and very informative. -Sarah

Course Description

- ✎ This course will describe how to appropriately choose a pain medication based upon individual patient and drug factors.
- ✎ Additionally, opioid medications will be evaluated in terms of risk versus benefit, with an emphasis on pain levels and the potential for addiction.
- ✎ Case anecdotes will include management of ocular pain, with specific emphasis on oral/systemic medications and how to protect both patient and practitioner.

Learning Objectives

- ☞ When given a patient case, choose an appropriate pain treatment plan for the management of ocular pain, in terms of drug choices based on pain level, dosing issues, and a monitoring plan for efficacy and toxicity.
- ☞ Identify and describe some of the potential signs, symptoms, and behaviors associated with opioid or substance abuse, and describe ways to respond to this issue.
- ☞ List systems available to evaluate a patient for potential opioid/substance abuse.
- ☞ Describe the treatment issues and options associated with the treatment of ocular pain in a patient with a drug abuse history.

Two major types of pain:


Nociceptive Pain – normal processing of stimuli that damages normal tissues; how pain becomes conscious;

- * responsive to non-opioids
 - * examples: NSAIDs, acetaminophen, steroids
- * responsive to opioids
 - * examples: codeine, hydrocodone, tramadol


Neuropathic: abnormal processing of sensory input by the peripheral or central nervous system;

- * treatment includes adjuvant analgesics
 - * sleep aids, nerve pain meds, muscle relaxers, anxiolytics

Nociceptive Pain- I was hit in the eye with a tree branch, and I think my eye is scratched



Nociceptive Pain- Healing Well



Pain is Complex and Confusing

Nociceptive Pain and Neuropathic Pain

February 9, 2022



Zoster

February 9, 2022



February 16, 2022



February 22, 2022



March 8, 2022



February 16, 2022





Not to Confused with

Neurotrophic

Stain Without Pain!


Actually, the OS is More Comfortable – What?




Corneal Sensitivity Testing



Cornea Sensitive Testing – Another Patient



Cornea Sensitive Testing – Yet Another Patient



What conditions have you used an Opioid?

🔗 Audience discussion



What Opioid do you prescribe?

🔗 Audience discussion



Drug Treatment Options...Neuropathic Pain

👁️ Why is this relevant?

👁️ Adjuvants – means “add on” medications

★ Some of them have addiction potential

📄 Anti-seizure medications that address nerve damage/inflammation

- MOA: work on the GABA system – similar to benzodiazepines (ex. Xanax)
- Gabapentin (Neurontin) – controlled substance in multiple states
- Pregabalin (Lyrica) – controlled substance in all 50 states

📄 Anti-anxiety and sleep medications

- Zolpidem (Ambien)
- Alprazolam (Xanax), Lorazepam (Ativan), Diazepam (Valium)

Pain Assessments and Scales

☞ Adds objective data to a patient's feeling of pain

- ★ It is a subjective problem to assess!

- ★ Remember...no patient should needlessly suffer!


☞ “Does the injury or wound or diagnosis fit the patient's presentation?”

- ★ It is important to be able to assess the degree of pain in a patient.



Combination Pain Scale...

PAIN MEASUREMENT SCALE



Drug Treatment Options...

Nociceptive Pain

3 Groups of analgesics

★ Non-opioids

- ☐ Acetaminophen (Tylenol)
- ☐ NSAIDs (Ibuprofen, naproxen sodium)
- ☐ Glucocorticosteroids (methylprednisolone, prednisone)

★ Opioids –

- ☐ Codeine (Tylenol with codeine)
- ☐ Hydrocodone (Vicodin)
- ☐ Tramadol (Ultram)

Controlled Substance Schedules

Schedule I – not considered to be medically necessary, research only

- ★ Heroin
- ★ “Medical” Marijuana
 - ☐ State control of marijuana and CBD
- ★ LSD
- ★ Mushrooms
- ★ Ecstasy

Schedule II – more likely to be abused (as compared to Schedule III, IV, V)

- ★ **Opioids, AKA “Narcotics”**
 - ☐ Oxycodone (OxyContin)
 - ☐ Hydrocodone (Vicodin, Lorcet, Norco)
 - ☐ Morphine (MSContin, MSIR)
 - ☐ Hydromorphone (Dilaudid)
 - ☐ Methadone
 - ☐ Fentanyl (Duragesic)

- ★ **ADD/ADHD meds:**
 - ☐ Methylphenidate (Ritalin)
 - ☐ Mixed amphetamine salts (Adderall)

Controlled Substance Schedules

Schedule III - Safer, less likely to be abused (as compared to Schedule II)

- ★ Combination products with APAP or ASA (codeine)
- ★ Esketamine – nasal spray for treatment resistant depression

Schedule IV – Safer, less likely to be abused (as compared to Schedule II and III)

- ★ Tramadol (Ultram)
- ★ Benzodiazepines (lorazepam, diazepam, oxazepam)
- ★ Sleep agents (zolpidem, etc.)

Schedule V – safest, least likely to be abused

- ★ Expectorants with codeine

Opioids “narcotics”

☞ Mainstay of therapy for the treatment of pain

☞ NO maximum daily dose limitation

☞ Useful for acute and chronic pain

Morphine Products

Morphine

★ Standard for comparison of other agents

☞ **MSIR** (IR caps) (q 3-4 hours prn)

☞ **MS Contin** (CR tabs) (q 8–12 hours)

☞ **Kadian** (CR caps) (q 12 – 24 hours)

☞ **Avinza** (CR caps) (q 24 hours)

Hydromorphone Products

Hydromorphone (Dilaudid) tablets – immediate release

Hydromorphone ER (Exalgo) tablets – extended release

 Used for severe pain

Codeine-Based

👁 Codeine – C3; Schedule III

👁 Hydrocodone – C2; Schedule II

👁 Oxycodone – C2; Schedule II

Codeine tablets

↳ WEAK analgesic: commonly used, so MOST have heard of it!

↳ Add acetaminophen/aspirin – Schedule III

★ **Tylenol #3** = 300 mg acetaminophen & 30 mg codeine

↳ Add expectorant – Schedule V

★ If you think someone won't try to get their hands on "codeine cough syrup" as a drug of abuse, you'd be surprised!!!

Oxycodone Products

Long-Acting, Extended-Release

OxyContin

Immediate Release; short-acting tablets

OxyIR (IR cap)

Roxicodone solution

with Acetaminophen:


Percocet and Endocet (oxycodone/APAP dose)

OxyCONtin (Controlled release tablets (q 12 hours...once in a while q 8 hours);
new formulation is out to help control abuse

Manual Crushing Followed by Dissolution



Crushed New Formulation




Crushed Original Formulation

Tampering for IV Abuse

- New formulation results in gelatinous material which cannot be drawn into a syringe for injection (the syringe is empty)

New formulation



Original formulation



Hydrocodone Products

Immediate-Release Products:

AS OF AUGUST 2014, hydrocodone products are ALL CII!!

Hydrocodone 7.5 mg + IBU 200 mg (Vicoprofen)

Hydrocodone + acetaminophen:

↳ “Vicodin” 5/300; 7.5/300; 10/300

↳ Lortab = 2.5/300, 5/300, 7.5/300, 10/300

↳ Norco = 5/325, 7.5/325, 10/325

Miscellaneous

👉 Fentanyl Patch (Duragesic)

- ★ MOST potent opioid
- ★ Black Box Warning against use in acute pain and in opioid naïve patients

👉 Methadone

- ★ Typically reserved for morphine/codeine allergic patients

Methadone tidbits...

☞ Chronic pain or opioid abuse deterrent

☞ 2-phase elimination

★ Alpha phase = 8 hrs

☑ Offers pain control

★ Beta phase = 16+ hrs

☑ Mitigates withdrawal symptoms

☞ Patient 1: On a short-acting pain med = likely being used to treat chronic pain

★ Twice per day dosing

☞ Patient 2: On methadone ONLY; lower doses

★ Once daily dosing

Tramadol

Tramadol (Ultram) tabs

Tramadol with 325 mg APAP (Ultracet), Tramadol ER tabs

- ★ Dual action: mu receptors & inhibits neuronal uptake of serotonin & norepinephrine
- ★ Lowers seizure threshold; increases serotonin levels
 - 📄 watch drug interactions with other meds that ↑ serotonin
 - Selective serotonin reuptake inhibitors (SSRIs): fluoxetine/Prozac
 - Migraine meds (“triptans”): sumatriptan/Imitrex
 - 📄 AS OF AUGUST 2014, NOW A C4 (Schedule IV)
 - 📄 “tramies” = abuse potential; helps decrease withdrawal symptoms

Opioid Allergies

🕒 If a patient states “codeine allergic”, ask appropriate questions...

★ “You have indicated that you have an allergy to codeine, can you describe what happens when you take codeine?”

- 📄 This is **SIGNIFICANT**, because if a patient is truly allergic to codeine, then they are most likely allergic to morphine, hydromorphone, oxycodone, hydrocodone, and tramadol
- 📄 **AND...**if they had an opioid IV after surgery, then their “reaction” may have been due to histamine release...
 - NOT always an allergic reaction

Opioid Allergies

👓 DO YOU KNOW WHAT A PATIENT CAN TAKE?

- 📄 Fentanyl
- 📄 Methadone
- 📄 Meperidine

👓 Assessing “allergies” appropriately helps practitioners sort through ACTUAL allergy potential and “placebo allergies”

- 📄 Fear versus drug seeking


Specific Medications Using Numeric Pain Scale

Mild pain = 1 – 3

- ☞ Acetaminophen (APAP; Tylenol)
- ☞ Ibuprofen (Advil, Motrin)
- ☞ Naproxen sodium (Aleve)
- ☞ Tramadol (Ultram) - low dose

Moderate pain = 4 – 6

- ☞ Tramadol (Ultram) – mid to high dosing
- ☞ Tylenol with codeine (Tylenol #3)
- ☞ Acetaminophen with oxycodone (Percocet)
- ☞ Acetaminophen with hydrocodone (Vicodin, etc.)



Specific Medications Using Numeric Pain Scale

Severe pain = 7 – 10


👉 Tylenol with hydrocodone (Vicodin, etc.) – higher doses

👉 Tylenol with oxycodone (Percocet, etc.) – higher doses

👉 Morphine (MSIR)

👉 Hydromorphone (Dilaudid)

👉 Fentanyl (Duragesic patch; Actiq lozenge on a stick)



Opioid Effects/ADRs

👁️ Sedation

👁️ Euphoria – mu receptors

👁️ Dysphoria/Hallucinations

👁️ Pruritis – allergy versus normal release of histamine

👁️ Nausea/vomiting

★ Triggers CTZ

★ Codeine “allergy”

Opioid Effects/ADRs

👁️ Confusion

👁️ Miosis

👁️ Respiratory depression – this is what kills a patient

★ *Mixing opioids with other CNS depressants*

- 📄 Alcohol
- 📄 Benzodiazepines
- 📄 Muscle relaxers
- 📄 Sleep agents
- 📄 Antihistamines
- 📄 Anti-seizure medications

Opioid Effects/ADRs


Withdrawal symptoms:

- ★ Short half-life agents are more likely to cause abrupt withdrawal symptoms
- ★ Sweating
- ★ High sympathetic tone: increase in heart rate and blood pressure, mydriasis
- ★ Agitation
- ★ Irritation
- ★ Irrational behavior
- ★ Symptoms disappear with (immediate) use of an opioid

Opioid Antagonists

Naloxone (Narcan) & Naltrexone (ReVia)

* Used to treat opioid overdose



Mixed Opioid Agonist-Antagonist

- ↳ Exhibit partial agonist or antagonist activity at the opioid receptors
- ↳ Agonist/Antagonist combinations for the TREATMENT of chronic pain
 - ★ NOT appropriate for the treatment of acute pain
 - ★ Morphine/Naltrexone (Embeda)
 - ★ Oxycodone/Naltrexone (Troxyca ER)
- ↳ Schedule II controlled substance

Mixed Opioid Agonist-Antagonist

↳ Exhibit partial agonist or antagonist activity at the opioid receptors

Agonist/Antagonist combinations for the TREATMENT of opioid abuse/addiction

↳ **Buprenorphine (Buprenex)**

↳ **Buprenorphine/Naloxone (Suboxone)**

↳ **Schedule III**

↳ **Adverse effects**

★ **Less respiratory depression & less abuse potential?**

↳ **Precipitate withdrawal in an opioid-dependent patient**

Painful Ocular Problems – things to consider...

👁️ Acute or chronic?

- ★ YOU are in charge!
- ★ Legal and ethical issues – do not allow yourself to be bullied by the patient!

👁️ Work with other practitioners!

👁️ Only a pain specialist should write RXs for CII medications for chronic pain issues

- ★ If something looks suspicious, then make inquiries! Especially before you write an RX for a drug that can be abused and/or sold!

Painful Ocular Problems – things to consider...

 Use the tools that are available!

- ★ State databases

 -  **PDMP** = Prescription Drug Monitoring Program

- ★ Pharmacists

Tolerance

👉 Escalation of dose to maintain effect (analgesia or euphoria)

★ Happens to everyone

👉 Regarding euphoria = may be life threatening because respiratory depression does not show much tolerance

“True Addiction” (formerly “psychological dependence”)

☞ Compulsive use despite harm

☞ Many times triggered by cravings in response to specific cues

- ★ Lifestyle is geared to the acquisition of the drugs
- ★ Borrowing from others, injecting oral formulations, prescription “loss”, requesting specific drugs (not always a sign...as some drugs just work better)

☞ Quality of life is not improved by the medication and eventually it becomes compulsive (“wanting without liking”)

☞ Relapse is very common even after “successful” withdrawal...it is a relapsing disease that is incredibly hard to treat

Identifying Behaviors of Abuse/Addiction

- 👁️ New patients that don't seem to "fit"
- 👁️ "fast talkers"
- 👁️ Strange allergies
- 👁️ Excuses for "loss" of meds or why they need "a strong pain medication"

Ways to respond

- 👉 Avoid getting “bullied”
- 👉 Avoid acting like you are judging the patient
- 👉 State data bases
 - ★ Call your local pharmacy/pharmacist
- 👉 Legal/ethical issues
 - ★ If you didn't write it down, then it didn't happen!
 - ★ If you accidentally give an addict a script for a pain medication, you won't get into “trouble”...

Substance abuse history...

☞ Avoid all opioids in a patient with a history of heroin use

- ★ This includes tramadol
- ★ May trigger dopamine reward and the drug “need”
- ★ Stick with higher doses of a NSAID +/- acetaminophen

☞ Patients with abuse history for other substances (ex. Benzodiazepines, alcohol, amphetamines)?

- ★ It is a judgement call
- ★ Some evidence to suggest that all addictive meds should be avoided!

Conditions Which May Require Pain Management

👁️ Large cornea abrasions

- ★ Cornea burn
- ★ PRK/PTK

👁️ Orbital trauma

👁️ Orbital blowout fractures

👁️ Scleritis





Optometric
Education
Consultants

Treatment of Pain Opioid Choices and Considerations

Greg Caldwell, OD, FAAO

Mid-Winter Getaway
Optometric Education Consultants

Sunday, January 28, 2024

